

BAD SPECIES

Henri Descimon & James Mallet

Laboratoire de Systématique évolutive
Université de Provence (case 5)
3 Place Victor Hugo
F-13331 MARSEILLE CEDEX 03, France
(descimon@up.univ-mrs.fr)
tel: 00-33 491-87-41-56
Fax: 491 10 63 03

Galton Laboratory, University College London,
Wolfson House, 4 Stephenson Way
LONDON NW1 2HE, England
(<http://abacus.gene.ucl.ac.uk/jim/>)
tel: +44-171-380-7412/7411
fax: +44-171-383-2048

Published in: Ecology of Butterflies in Europe (eds. Settele, J., Konvicka, M., Shreeve, T., & Van Dyck, H.). Cambridge University Press. pp. 219-249.

N.B. This version differs somewhat from the published version, lacking some of the corrections in that document, but also containing some updates that were not incorporated into the final version. JM Jan 2010

I. Introduction: species concepts and taxonomic practice

Taxonomists, when describing a new species, often added the term *bona species* after the Linnaean binomial. The implication is that there are also *malae species*. A “bad species” is a taxonomic unit that misbehaves with respect to criteria used to delimit species. There are a wide array of species definitions linked to theories of speciation and evolution (Harrison, 1998; Coyne & Orr, 2004) and there have been many debates, which often become abstruse and epistemological (Wilson, 1999, Hey, 2006). The biological species concept (BSC), based on reproductive isolation and associated with the theory of allopatric speciation, prevailed for many years. More recently, the advent of numerical taxonomy (Sokal & Crovello, 1970) and cladistics (Hennig, 1968) has upset the earlier certainty. The establishment of a basis for conceiving (Maynard-Smith, 1966) and observing (Bush, 1969) sympatric speciation led to suspicions that species were more indefinite, even locally, than architects of the modern synthesis had imagined. Today, two different consensus seem to be building among evolutionary biologists. The species concept either takes the form of a minimal, Darwinian, definition which is agnostic about evolutionary mechanisms to allow universal applicability (Mallet, 1995; Feder, 1998; Jiggins & Mallet, 2000), or attempts to combine a variety of species concepts together (Templeton, 1998; de Queiroz, 1998; Coyne & Orr, 2004). Under both views, species may evolve or be maintained via multiple different routes.

Species concepts and criteria: speciation theory and systematic practice

When treating an actual fauna or flora, the central problem is of the purely taxonomic criteria for species status. For a long time, four kinds of criteria have been used to group members of a species: character-based or “syndiagnostic” criteria (which may use morphological or genetic traits); phylogenetic or “synepigonic” criteria; reproductive, “mixiological”, or “syngamy” criteria; and finally geographic criteria, particularly “sympatry,” “cohabitation,” or geographic overlap (Poulton, 1904; see also Jordan, 1905; Rothschild & Jordan, 1906; Cuénot, 1936). To be distinct at the level of species, taxa should provide at least some of these four kinds of evidence. With the advent of biological species concept (BSC; Dobzhansky, 1937; Mayr, 1942), the main emphasis was put on reproductive isolation (i.e. “mixiological”) criteria. This caused something of a divorce between evolutionary theory and taxonomic practice. On the one hand, overwhelming amounts of work has been carried out on the genetics and evolution of species: studies of genetic structure within species, interspecific crosses in the laboratory, and field studies on hybrid zones (Barton & Hewitt, 1989; Berlocher, 1998; Coyne & Orr, 2004). On the other hand, practising taxonomists often continue to use “syndiagnostic” methods based mainly on morphological characters.

Indeed, when taxonomists have a sample of specimens coming from an unexplored geographic area, they can find morphological differences with taxa already described, but it is difficult to determine whether they are due to a few pleiotropic gene changes (i.e. the new samples are merely morphs of described taxa), to intraspecific geographic variation (subspecies), or to differentiation at full species level. Sometimes, rare hybrids between well-known species have even been mistaken for “good” species. Since they are inaccessible, other criteria are simply ignored. Although they can reveal much about mixiological criteria, chromosomal and molecular characters are often used in much the same way as early taxonomists used morphological data; for instance, differences in chromosome numbers, or the presence

of diagnostic allozyme loci has been considered proof of distinct species, without consideration of geography or genetic relationships. We argue that these biological characteristics cannot be ignored.

Study of ecological niches is particularly important for associating morphological or genetic differences with different habitats (Sneath & Sokal, 1973). Mayr, in later versions of his BSC (1982) argued that each species “occupies a biological niche in nature”. Adaptive evolution is recognized as a primary means of both splitting and maintenance of separate lineages (Van Valen, 1976; Templeton, 1989, 1994, 1998; Andersson, 1990; Baum & Larson, 1991; Schluter, 2000). Sympatric speciation also involves ecological differentiation (Bush, 1969, Feder, 1998), and increasing evidence suggests that ecological divergence may directly cause reproductive isolation (Dodd, 1989, Schluter, 2001).

Nonetheless, mixiological criteria remain the most important within the BSC conceptual framework. They are reached through observation of the relations between the taxa either in sympatry, or in hybrid zones in the case of parapatry (Jiggins & Mallet, 2000) – the latter are considered as “natural laboratories for evolutionary studies” (Hewitt, 1988). Modelling as well as empirical studies suggest that hybrid zones can act as a barrier to gene flow (Barton & Hewitt, 1989). Within them, the intensity of hybridization may vary. If hybrid genotypes predominate, the hybrid zone is considered “unimodal”, while, if genotypes are predominantly parental, with few intermediates, it appears phenotypically “bimodal” (Harrison & Bogdanowicz, 1997; Jiggins & Mallet, 2000). Pairs of species that cohabit broadly and hybridize regularly can be studied genetically in the same way. In hybrid zones, the mixiological criterion of species depends on the fraction of genes that are actually exchanged between the taxa. Hybrids can be detected using morphological criteria, but this can be inaccurate, which makes it hard to estimate gene flow. Gene exchange, or introgression (Stebbins, 1959), may transfer important genetic variation in some cases of adaptive evolution, especially in plants (Arnold, 1992, 1997; Mallet, 2005) In birds and fish, hybridization is widespread (Grant & Grant, 1992) and may be involved in rapid adaptive radiation and speciation (Grant & Grant, 1998; Seehausen, 2003). This also seems likely in *Heliconius* butterflies (Gilbert, 2003; Bull *et al.*, 2006). Introgression can affect the mitochondrial genome (Aubert & Solignac, 1990) but, in Lepidoptera, where the Y-bearing sex is the female, Haldane’s rule severely hinders mitochondrial introgression (see below and Sperling, 1990, 1993; Aubert *et al.*, 1997).

Based on the ideas of Mallet (1995) and Feder (1998), the separation of gene pools during speciation has been dubbed “the genic view of speciation” by Wu (2001): speciation may not take place *via* separation of the whole gene pools, as postulated by the Dobzhansky-Mayr theory of speciation, but initially concerns only genes actively involved in reproductive isolation. The rest of the genome may still undergo sufficient gene flow to prevent differentiation, except in genomic regions tightly linked to “speciation genes” (Ting *et al.*, 2000). But what are speciation genes? Genes involved in divergent adaptation and mate choice should diverge first, and those causing hybrid sterility and inviability should be expected to diverge only after initial genetic separation. Complete separation should result from reinforcement of sexual isolation and further ecological differentiation (Noor, 1999). Although Wu’s genic view of speciation elicited an immediate rebuttal from the father of the BSC (Mayr, 2001), it is clear that the proposed scheme is not that different from the “classical” view of speciation according to Mayr. The most important distinction is that Wu’s modification of Mayr’s speciation scheme renders it compatible with a more

substantial phase of gradual divergence in sympatry or parapatry.

An array of varied data obtained from difficult or “bad” taxa can be used to support or refute the presence of additional species within a sample. The more concordant the data are, and the more bimodal the frequency distributions of phenotypes and genotypes, the more likely separate species status will be granted. These are methods termed “genealogical concordance” or “genotypic clustering” (Avice & Ball, 1990; Mallet, 1995). Similar “syndiagnostic” procedures were, in fact, being applied to morphological characters long before Darwinian times (Adanson, 1763). As early as 1930, Nilsson (cited by Cuénot, 1936) used the term “genotypenkreis” to characterise species in *Salix*, a plant genus prone to hybridization.

This ideal procedure for species delimitation, careful study in zones of contact, is not always possible. In cases where concordance between criteria is imperfect, some argue for distinction at species level, and others against it. For instance, cryptic or sibling species (Dobzhansky, 1937; Mayr, 1963) fail to show diagnostic morphological characters; species that are otherwise well-characterized apparently share the same ecological niche; hybrid zones can be unimodal in some areas and bimodal in other parts of the range. Molecular markers may be strongly differentiated among populations within species; in other groups, species clearly distinct using other criteria can show little molecular differentiation, especially if speciation is recent compared with the rate of molecular divergence.

Cohabitation: the lumpers' species criterion adopted here

The touchstone of all criteria for separate, biological species is the test of “cohabitation”: whether overlapping populations produce unimodal (in which case subspecies might be designated), or bimodal (in the case of separate species) morphological and genotypic frequency distributions. This procedure dates from the late 19th Century, and was promoted particularly vigorously for the Lepidoptera by Karl Jordan (e.g. Jordan & Rothschild, 1906). Other species criteria that do not depend on degree of hybridisation or intermediacy in areas of overlap are also in use today. In particular, Cracraft’s (1983, 1989) “phylogenetic” or “diagnostic” concept is contributing to taxonomic inflation of “species” numbers in birds, primates, and other taxa (Isaac et al., 2004), even when no new populations have been discovered. In butterflies, the prohibitive diversity of morphologically or genetically diagnosable local populations, usually referred to in our literature as “subspecies,” has tended to prevent such rampant splitting (for the moment). Here, we adopt this traditional and more inclusive, polytypic or “lumpers'” criterion for species.

When sympatric taxa hybridize very rarely, they can be classified as separate species. But what can be concluded if the units to be compared are not in contact? Breeding and crossing experiments provide an apparent solution, but this can be misleading. In particular, viability of hybrids in the laboratory may appear normal while, in nature, hybrids could be severely disadvantaged. Pre-mating barriers to hybridisation can also be reduced under artificial conditions. In both cases, the degree of mixiological separation estimated can be spurious.

Whenever there is conflict between criteria, or whenever regular hybridisation occurs, in spite of the fact that the taxa remain to some extent morphologically, ecologically or genetically distinct, or if populations are allopatric but seem at that stage of divergence at which species fusion is doubtful, one may speak of “bad species”. The tools used in making a decision on the rank of taxa at this stage of divergence include morphological, chromosomal (karyological), molecular, and

ecological characters. In addition, one may cross such taxa, to obtain criteria relevant to reproductive isolation and introgression, keeping in mind the *caveat* previously invoked. These tools are described in detail in the appendix.

As with any term, "species" must have a definition that depends partly on theoretical considerations. At this point, one might ask two main groups of questions: 1- Do species exist as real entities in nature? Or are they a construct of the human desire for categorization and classification? 2- What are species made of? How do they arise? How are they maintained? And are species a homogenous rank from this evolutionary point of view? To answer such questions, it is necessary to investigate actual problem cases in some depth, which is the main aim of the rest of this chapter.

II. How common are bad species in European butterflies?

It is often said that, although there are disagreements about species concepts, there are few cases where our ability to delimit species is severely challenged (e.g. Mayr, 1963). However, hybridization and bad species are rather more common than field guides tend to mention. Taxonomists overlook "dubious" individuals (which may often be hybrids) because they make species discrimination more difficult. Natural hybridization occurs between around 10% of all animal species, although there are many groups where hybridization rates are greater (Mallet, 2005). Here we provide collated data on European species, one of the best studied faunas in the world (Table 1). Overall, around 16% of the 440 butterfly species are known to hybridize with at least one other species in the wild. Of these perhaps half or more are fertile, and show evidence of backcrossing in nature.

III. Case studies: the practice of European butterfly taxonomists at species level.

European butterflies are taxonomically well known. In the first comprehensive work on European butterflies, Higgins and Riley (1970) enumerated 371 species (including the skippers); in a recent book of the same scope, Tolman & Lewington (1997) record 440 species, 69 more. Amongst the "new" European species, hardly any are actually new finds; many arise from "taxonomic inflation", the upgrading of previously known subspecies to species level, or discoveries of known non-European species just inside the boundary (Dennis, 1997; Isaac *et al.*, 2004). In this section, we present an analysis of some decisions which illustrate how splitting and/or lumping has been performed in particular cases.

1. The genus *Hipparchia*: splitters and lumpers at work.

Some genera have undergone especially intense splitting, like the graylings (*Hipparchia* and *Neohipparchia*). According to Higgins & Riley (1970), there were only 10 species in Europe. Today, there are 19 (Tolman & Lewington, 1997), to which one more, *H. genava* can be added according to Leraut (1990). Mostly, this proliferation is due to elevation to species rank of forms inhabiting islands or other disjunct geographic regions (e. g. *H. azorina*, *H. caldeirensis* and *H. miguelensis* in the Azores). However, this is not true for *H. alcyone* and *H. genava*, between which Leraut records a hybrid zone. In a revision of the genus (Kudrna, 1977) elevation to species rank was based only on morphology. Morphometric analyses of multiple, well-replicated samples in the *semele* group based on genitalia, wing pattern measurements and allozyme electrophoresis were later carried out by Cesaroni *et al.* (1994), who showed convincing congruence between the morphometric analysis of genitalia and allozymes, although wing patterns followed an obviously different

evolutionary pathway. The number of taxa with specific status was reduced by Cesaroni *et al.* from eight to five. As the taxa were largely allopatric and often insular in distribution, cohabitation and hybrid zone criteria cannot be tested. Assignment to species level was therefore performed on the basis of “sufficient” genetic distance (Nei’s *D* between 0.07 and 0.26).

Later, Jutzeler *et al.* (1997) presented another treatment of the same group. Although devoted mainly to meticulous morphological description of certain taxa and their first instars, and lavishly illustrated with SEM pictures and excellent colour plates, the specific status of the various taxa was also discussed. The authors, it turns out, are extreme “splitters”, and even cite Cesaroni *et al.* (1994) to justify splitting – in complete contradiction to that paper. No morphometric analyses were performed while making these controversial decisions. More recently, even more “insular splitting” has been carried out by Jutzeler *et al.* (2003a, b): taxa from the Tyrrhenian islands were raised to species on the basis of morphological and bionomic differences with continental relatives, again without any morphometric, karyological, mixiological or molecular justification. Most of these “new” species are allopatric. We tend to side with the more conservative views of Cesaroni *et al.* (1994).

2. *Agrodiaetus admetus* and the “anomalous blue” group: chromosome variation and allopatry.

According to Lukhtanov *et al.* (2003), “this complex is a real stumbling block in the taxonomy of the genus [*Agrodiaetus*]”. In a careful study using the “classical” tools of typological taxonomy, Forster (1956) was uncertain about the taxonomic status of only a few forms or “bad species”. Soon thereafter, de Lesse (1960a) used karyology to show that the picture was not simple but death prevented him from carrying his work further. The *admetus* group of *Agrodiaetus*, which included only three species in Higgins and Riley (1970), was raised to nine some 35 years later (Tolman & Lewington, 1997; Wiemers, 2003).

In *Agrodiaetus*, the males are generally blue, but the “anomalous blues” all have similar, chocolate-brown uppersides in both sexes. In 1970, the species recognized were *A. admetus*, ranging from Eastern Europe to Asia Minor, *A. fabressei* known only from Spain, and *A. ripartii* from scattered locations from Spain to Asia Minor. This treatment was supported by karyotyping: $n=78-80$ for *admetus*, $n=90$ with two large unequal chromosomes for *ripartii*, and $n=90$, with two large and two medium-sized chromosomes for *fabressei* (de Lesse, 1960a). The taxa *fabressei* and *ripartii* cohabited without admixture in some Spanish localities (de Lesse, 1961a).

The situation became more complex when wide karyotypic variation was found in Turkey and later in parts of Europe. Below, we present the taxa recognized by various authors (Hesselbarth *et al.*, 1995; Eckweiler & Häuser, 1997; Häuser & Eckweiler, 1997; Carbonell, 2001; Lukhtanov & Dantchenko, 2002a,b, 2003; Wiemers, 2003; Kandul *et al.*, 2004). Taxa with no information on chromosome number are omitted, as are taxa of obviously subspecific rank:

Table 2.

Species of <i>Agrodiaetus</i> (according to Wiemers, 2003; Tolman & Lewington, 1997)	Distribution	Chromosome number (<i>n</i>)
<i>admetus</i> Esper	Bulgaria	80
” ”	Turkey	78-80
<i>alcestis</i> Zerny	Lebanon	20-21
<i>aroaniensis</i> Brown	Peloponnese	48
<i>dantchenkoi</i> Lukhtanov <i>et al.</i>	Turkey	42
<i>demavendi</i> Pfeffer	Iran, Turkey	68-71
<i>eriwanensis</i> Forster	Armenia	32-34
<i>fabressei</i> Oberthür	Spain	90 (86+2+2)
<i>galloi</i> Balletto & Toso	S. Italy	66
<i>humedasmae</i> Toso & Balletto	N. Italy	38
<i>interjectus</i> de Lesse	Turkey	29-32
<i>karacetinae</i> Lukhtanov & Dantchenko	Turkey	19
<i>nephoiptamenos</i> Brown & Coutsis	N. Greece	8-11, or ~90*
<i>ripartii</i> Freyer	Spain-Turkey	90 (88+1+1)

* There are contradictory numbers counted by Brown & Coutsis (1978) and de Prins (unpublished); the $n=90$ estimate seems most likely (Wiemers, 2003).

More recently, allozyme studies have cast doubt on this multiplicity of species. *A. ripartii*, the most widespread, proved as homogeneous genetically as in its karyotype; this is also true, to a lesser degree, for *A. admetus*. *Agrodiaetus fabressei* and the other taxa are poorly resolved and there is little correlation between allozymes and karyotype (Mensi *et al.*, 1994). More recently, mitochondrial and nuclear DNA sequencing suggest that “brown” *Agrodiaetus* are polyphyletic. The wing colour switch from the “primitive” blue colour to brown in males seems to have occurred twice: once in the “*admetus*” group and once in *fabressei* (Wiemers 2003; Kandul *et al.*, 2004). Most distinguishable entities are allopatric, and the only exceptions are the aforementioned *A. fabressei* and *A. ripartii*, and four species found close together in the Turkish Van province (Lukhtanov *et al.*, 2003). In most other cases, nobody knows what would occur if these genetic entities flew together.

Clues are provided by the *fabressei-ripartii* case, which have the same chromosome number, but differ in details of the karyotype. They comply with the cohabitation criterion and are genetically distant (Lattes *et al.*, 1994). Clearly, there is little doubt that these are good (albeit sibling) species. However, they are almost impossible to identify using morphology where they co-occur, since neither wing pattern nor skeletal morphology provide reliable criteria: karyotype and DNA sequencing is virtually the only way to assure identification (Lukhtanov *et al.*, 2003).

Chromosomal information has also been used by Munguira *et al.* (1994), who merged the Spanish *agenjoi* Forster and *violetae* Gomez-Bustillo *et al.* into known species, *fabressei* and *ripartii*.

Chromosome structure is unstable in *Agrodiaetus* and rearrangements are common even within populations, leading to the formation of multivalents during meiosis (Lukhtanov & Dantchenko, 2002a, b; Lukhtanov *et al.*, 2003). Limited abnormalities seem not to affect viability, although selection should eliminate most rearrangement polymorphism eventually. Why is chromosome structure so unstable in *Agrodiaetus*? Kandul *et al.* (2004) argue that tolerance of chromosomal polymorphism is related to centromere structure, and suggest that destabilization of chromosome numbers may be due to locally abundant transposons. In allopatric populations of *Agrodiaetus*, elimination of differences will not take place and the karyotype diverges rapidly until a point of no return is reached, giving rise to a great deal of geographic variation, and ultimately speciation. Similarly, Wiemers (2003) boldly states that “changes in the number of chromosomes do not lead to sympatric speciation, but instead appear as a by-product of allopatric speciation and such young species could only occur in sympatry after a sufficient differentiation in their phenotype to exclude erroneous matings”.

3. *Leptidea sinapis* and *L. reali*: sibling species and the almost “perfect crime,” with a comparison to the situation in *Melitaea athalia*.

Until the end of the 20th century, nobody suspected that two separate species lurked within the wood white, *Leptidea sinapis*. In 1962, Réal noticed that two different seasonal forms flew together in the French Eastern Pyrénées, without considering the possibility that two species were involved. By the late 1980s, after morphological studies on the genitalia, Lorković suggested to Réal that there were indeed two species. The latter described a new species under the name *lorkovicii* in 1988, an invalid name replaced by *reali* (Reissinger, 1989). Further study confirmed that the two forms, characterized by male and female genitalia, were distinguishable and sympatric across much of Europe (Lorković, 1993; Mazel & Leestmans, 1996); in particular, the penis is short in *sinapis*, and long in *reali*. There are correlated differences in the females, with short vs. long *ductus bursae*. This strongly suggests a “lock and key” mechanism is involved. Although other barriers may be present, it seems likely that these differences can explain reproductive isolation between the taxa. In contrast, earlier attempts to find reliable differences in wing pattern and ecology were in vain. *L. sinapis* is present everywhere in Western Europe, while *reali*, if present, is always in sympatry with it.

Although the existence of two “good” species is likely, it could be argued that there is merely a genitalic polymorphism, similar to that in *Melitaea athalia* and *M. celadussa* (see below). To address this point, a study based on multivariate morphometrics of genitalia, allozymes and mtDNA sequencing was undertaken by Martin *et al.* (2003) on 6 populations from Southern France. A 728 bp fragment of the ND1 gene showed a reliable and constant 3% divergence between the entities. Among 16 enzyme loci, none was completely diagnostic, but *Ak* and *Pgi* showed highly significant differentiation. Multivariate analysis demonstrated two well separated “genotypic clusters”, with strong linkage disequilibria between loci. Furthermore, allozymes and the mtDNA were concordant. Morphometrics carried out on genitalia also yielded good concordance with molecular data, although there was some (< 5%) overlap between the taxa. In 163 individuals of the two species, no hybrid was detected; the few individuals with doubtful genitalic measurements were

clearly assigned to one or other taxon by molecular markers.

The necessity of dissecting individuals for identification makes ecological study difficult, and it was at first thought that the species fly together and share most foodplants. This should contradict Gause's principle but could explain the lack of consistent differences in wing pattern. However, the population genetic structure of the two species is somewhat different: *L. reali* is less polymorphic at allozymes (with heterozygosity $0.09 < H < 0.14$ in *sinapis* and $0.05 < H < 0.07$ in *reali*: Martin *et al.*, 2003). Females and, to a lesser extent, males of both species discriminate between the species during mate choice, and only intraspecific matings occurred in captivity (Freese and Fiedler, 2002). The two species are now known to differ in ecology: *L. sinapis* is a widespread generalist on various herbaceous Leguminosae from both wet and dry habitats, while *L. reali* specializes on *Lathyrus pratensis*, a plant confined to moist grasslands. In 347 localities in the Drôme department (southern France) where *L. sinapis* and/or *L. reali* were observed, *L. sinapis* was alone in 55% of the study sites, and *L. reali* in 22%, whereas both species were found together in 23% of them (Amiet, 2004). There are also differences in phenology, response to temperature and habitat choice (Friberg *et al.*, 2007). The situation seems to reverse in Eastern Europe, where *L. sinapis* becomes confined to warmer areas (Benes *et al.* 2003). Freese and Fiedler (2002), in their mainly laboratory-based study, concluded that “the two species are only weakly differentiated in ecological terms;” indeed, their egg-laying tests showed only a weak preference for *L. pratensis* in the females of *L. reali*; the larvae of both species prefer, and perform better on another legume, *Lotus corniculatus*, a result rather discrepant with Amiet's (2004) field observations.

As in almost all “perfect crimes”, once the first clue was discovered, a cascade of confirmatory data was quickly revealed. At the end of the 19th century, the earliest dissectors of genitalia, such as Reverdin, could well have studied a series of *Leptidea* male genitalia and discovered the two species.

The latter did just this with *Melitaea athalia* (Reverdin, 1920, 1922), where two types of male genitalia were associated with two biogeographical entities, and he therefore split them into separate species. However, later study showed that the morphology of male genitalia was unimodal within a hybrid zone between the two taxa. The width of the hybrid zone varied from a few to several tens of kilometres (Bourgogne, 1953). Since this differentiation is not associated with large and constant differences in allozymes or mtDNA, as in *Leptidea* (Zimmermann, unpublished), in *Melitaea*, species separation was premature.

4. *Zerynthia rumina* and *Z. polyxena*: relativity of mixiological criteria

The genus *Zerynthia* contains two species, both recognized since the dawn of entomology, *Z. rumina*, a Western Mediterranean species, and *Z. polyxena*, Eastern Mediterranean. They overlap in Southern France, where they display marked ecological differentiation, while in areas where only one species is found, both have a more extensive niche. Besides wing pattern differences, there are diagnostic alleles between, with Nei's $D \approx 0.80$, (Braconnot, unpublished) and strong divergence in mitochondrial and nuclear gene sequences (Nazari *et al.*, 2007) – there is no doubt they are “good” species! Both display marked intraspecific differentiation: wing patterns of the French subspecies *Z. rumina medesicaste* and *Z. polyxena cassandra* clearly differ by wing pattern from their respective nominal subspecies, but variation forms a wide cline within a continuous distribution.

Natural hybrids between the species are scarce (only five are known to HD), but interspecific pairing has been observed in the field (de Puységur, 1947). A large series of crosses within and between species has been performed by HD, although only some have been published (Descimon & Michel, 1989). When *Z. rumina medesicaste* was crossed with *Z. r. rumina*, remarkable hybrid vigour was observed in the F1, followed by strong hybrid breakdown in the F2 (i.e. F1 x F1): arrested embryonic development, larval weakness, and difficulties of pupation. Less than 5% of ova reached the adult stage in about 10 parallel broods. The low viability persisted in further crosses; only backcrosses, with either parent subspecies (or, paradoxically, with *Z. polyxena*), restored viability. Crosses between *Z. p. polyxena* from Greece and *Z. p. cassandra* from southern France also produced F1 hybrid vigour, and some F2 (F1 x F1) hybrid breakdown. However, the F2 viability was not too low (around 25%), and further crosses (F2 x F2 and more) displayed markedly enhanced viability: incompatibility therefore seemed less marked than in the first case. Crosses between Austrian and French *Z. polyxena* produced no F2 hybrid breakdown.

Mate choice was studied in cages containing 10 males and 10 females of each species. Only intraspecific matings were observed (including the aforementioned distinct subspecies), demonstrating strong prezygotic barriers between species. All females proved to have mated, and one female *polyxena* produced offspring consisting partly of *polyxena* and partly of hybrids. Clearly, she had mated twice, and with males of each species. The hybrids were viable, but while the F1 x F1 cross resulted in no offspring, backcrosses with *polyxena* and *rumina* were successful. The backcross hybrids from either side could, however, be crossed with the more distant parental strains. Thus backcrossed individuals, which had 3/4 of their genes from one species and 1/4 from the other, gave symmetrical F3 progeny with 3/8 *rumina* - 5/8 *polyxena* offspring and the reciprocal; the same scheme was applied in F4 and beyond. The possibilities for complex crosses increased with the rank of hybridization and some were practised (for a complete account, see Descimon & Michel, 1989). The hybrids were viable provided they had at least one complete unrecombined genome from a parental strain. Much more surprisingly, two later hybrid x hybrid crosses (not many were tried) gave fairly viable offspring, with no significant departures from 1:1 sex ratio or diapause abnormalities. In spite of strong pre-mating isolation between the pure species, female hybrids were attractive to males of either species, and male hybrids were attracted to any female. Similar results on hybrid sexual attractiveness have been obtained in a number of other butterfly species; e.g. *Heliconius* (McMillan et al., 1997; Naisbit et al., 2001).

It was not possible to continue the crosses, but some clear facts emerge. First, F2 hybrid breakdown is not absolute in interspecific crosses; second, it is not limited to interspecific crosses; it may take place between subspecies, as is known in other species (e.g. Oliver, 1972, 1978; Jiggins *et al.*, 2001). The latter fact is particularly paradoxical, since, within both species, broad, clinal, unimodal hybrid zones connect “incompatible” populations. Careful field work could well disclose interesting features in these contacts. Hybrid inviability is therefore probably not a useful species criterion on its own in crosses between geographically distant taxa. The ease of playing ping-pong with the two species once initial barriers have been ruptured shows that there is no absolute threshold of post-zygotic incompatibility at the species level.

5. Frequent hybridization and introgression in sympatric swallowtails: *Papilio machaon* and *P. hospiton*; *Parnassius apollo* and *P. phoebus*

a. The swallowtail and Corsican swallowtail

Hybridization is widespread in swallowtails, especially in N. America (Sperling, 1990). Hybrids between the Eurasian *Papilio machaon* and the endemic *P. hospiton* of Corsica and Sardinia have been known for a long time (e.g. Verity, 1913). Although their habitats and distribution in Corsica are very different, there is a frequent overlap, and hybridization occurs regularly. Crosses revealed two especially important postzygotic barriers (Clarke & Sheppard, 1953, 1955, 1956; Clarke and Larsen, 1986):

1. An almost total inviability of F1 x F1 hybrid crosses, originally mistaken for F1 sterility. However, non-hatching ova were not “sterile”; instead embryos show arrested development at various stages between early segmented embryos and fully-developed larvae unable to break out of their egg shell.

2. Strong Haldane’s rule F1 hybrid effects. In *hospiton* male x *machaon* female crosses reared in Britain, female hybrid pupae became “perpetual nymphs”, that is pupae which are unable to resume development. However, in other *Papilio* interspecific hybrids with extended diapause, ecdysone and insulin injections can trigger development (Clarke *et al.*, 1970; Arpagaus, 1987). Descimon and Michel (in Aubert *et al.*, 1997) showed that insulin could also trigger development in *machaon* x *hospiton* hybrids.

Both reciprocal F1 crosses and various backcrosses proved possible. The experiments were carried out in the Paris region, in an oceanic climate, and in Marseilles, on the Mediterranean, but under long photoperiod summer in both cases (Aubert *et al.*, 1997). In the case of *hospiton* male x *machaon* female crosses, results depended on rearing conditions. In Paris growth and developmental time of males was normal, but the female pupae, which were markedly bigger than those of either parental species, became “perpetual nymphs”, as found by Clarke and Sheppard (1953). In Marseilles, females did not enter diapause and gave large, viable females. The other possible F1 (*hospiton* female x *machaon* male) again gave healthy hybrid males, but females were small, with accelerated development and no diapause, in both towns. F1xF1 crosses gave almost complete inviability at various stages of early development, as before. On the other hand, backcrosses were all viable. F1 hybrid females, in particular, appeared not to be sterile, whether they had *hospiton* or *machaon* as mothers.

The results suggest that introgression is possible. Allozyme and RFLP (Restriction Fragment Length Polymorphism) analysis of mtDNA markers show strong differentiation between the two species, with diagnostic alleles at some loci and a rather high Nei’s *D* and mtDNA sequence divergence (Aubert *et al.*, 1997; Cianchi *et al.*, 2003). Putative hybrids found in different localities in Corsica and Sardinia were most probably F1s, and from both reciprocal crosses. No individuals were found with introgressed mtDNA RFLP types in a large sample, suggesting a lack of mitochondrial introgression. However, the same was not true for nuclear loci. Alleles from *hospiton* were found in Corsican *machaon*, but were always absent in continental *machaon* (Aubert *et al.*, 1997; Cianchi *et al.*, 2003). The frequency of hybrids was lower in the Italian than the French dataset (approx. 1% vs. 5%), but this is probably because HD collected especially avidly in areas of cohabitation, whereas many samples by the Italians contained only one species.

Classically, *hospiton* is considered single-brooded, while *machaon* is multi-brooded. However, broods reared from from wild Corsican *hospiton* females give a

proportion (5-100%) of non-diapausing pupae (Aubert *et al.*, 1996a). Diapause control in *P. hospiton* (and in *P. machaon*) is highly heritable but not simple; temperature and photoperiod act in combination, with threshold effects which interact strongly with genetic factors. Multi-brooded individuals are particularly common where *hospiton* feeds on *Peucedanum paniculatum*, a perennial evergreen umbellifer endemic to northern Corsica; this plant is suitable throughout the warm season. Observations in July and August confirm the existence of the second brood (Aubert *et al.*, 1996a; Guyot, 2002; Manil & Diringier, 2003). In most regions of Corsica and throughout Sardinia, the main foodplant, *Ferula communis*, withers down as early as May onwards. Even here, late larvae can be found when roadside mowing during late summer renders resprouting *Ferula* available (Descimon, pers. obs.).

Aubert *et al.* (1997) suggest that multi-voltinism in *P. hospiton* may result from introgression from *P. machaon*. This hypothesis was criticized by Cianchi *et al.* (2003) because of doubt in the existence of the second brood of *P. hospiton* (this argument is not tenable, as we have seen). Of more weight is the difficulty of distinguishing ancestral from introgressed polymorphisms. Nonetheless, Cianchi *et al.* (2003) found up to 43% *hospiton* allozymes in *machaon* on the islands, though never present on the mainland, and they argued that this was due to introgression. Conversely, they found only a scattering of *machaon* alleles in *hospiton*. They argued that this introgression was mostly ancient and that reinforcement of interspecific barriers took place early during the secondary contact. This conforms to the commonsense prediction that what we observe today is an equilibrium between gene flow and selection against introgression (Descimon *et al.*, 1989).

b. The Apollo (*Parnassius apollo*) and small Apollo (*P. phoebus*)

The Apollo is a montane butterfly, widespread from Altai in central Asia to the Sierra Nevada in southern Spain. The small Apollo has a more restricted, higher elevation distribution; in Europe, it occurs and can hybridize with the Apollo only in the Alps. The species always occur in close proximity (dry, sunny slopes for *P. apollo* and banks of torrents and rills for *P. phoebus*), but this does not ensure hybridization. Not only are their preferred flight environments different, but *P. phoebus* also flies earlier in the year. Therefore, it is only in localities where the two kinds of habitats are closely interspersed and phenology is perturbed that hybridization takes place, often at rather high frequency (Descimon *et al.*, 1989). In some localities, hybrids are observed almost yearly; in others, they occur only following a snowy winter, when avalanches accumulate in the bottom of thalwegs. Thus, rather “soft” pre-mating barriers, such as habitat and phenology differences prevent hybridization. In captivity, mating between male *apollo* and female *phoebus* is often observed, and hand-pairing easy. The reverse cross is more difficult, due to the small size of male *phoebus*. F1 hybrids display typical vigour and females are not perturbed in diapause (which takes place in the first larval instar, inside the egg shell). Field observations on wild hybrids show a strikingly perturbed behaviour: males fly restlessly, constantly roaming between the types of habitat preferred by both parent species. In captivity, male hybrids backcross freely with females of both species and are highly fertile, but female hybrids are inevitably sterile, producing numerous small ova that never hatch.

Morphometric analyses of natural populations strongly suggested backcrossing as well as F1 hybrids in the field (Descimon *et al.*, 1989). Using four diagnostic allozymes and several other loci with different allele frequencies in the two species, F1 hybrids and backcrosses were detected (Descimon & Geiger, 1988). One individual with the pure *apollo* wing pattern was heterozygous at one of the diagnostic

loci, suggesting that backcrossing continues beyond the F2. Mitochondrial DNA analysis showed that hybridization took place in both reciprocal directions but also that backcrossing could involve hybrid females (Deschamps-Cottin *et al.*, 2000). While this contradicts findings from some captive broods (Descimon *et al.* 1989), it conforms to others (Eisner, 1966). Once again, introgression in nature seems possible and is demonstrated by the field results.

c. Comparisons between the two hybridizing pairs of papilionids

It seems clear that most would regard the four swallowtails treated here as four distinct, if somewhat bad species. They are readily distinguishable on the basis of morphology, allozymes, and mtDNA. Allozyme and mitochondrial divergences suggest an age of around 6 Myr for the *Papilio machaon*-*P. hospiton* pair (Aubert *et al.*, 1999), and a similar age is probable for *Parnassius apollo* and *P. phoebus*. Regular hybridization is therefore not necessarily a sign of incomplete speciation, but rather of the inability of the taxa to erect complete pre-mating barriers.

In conclusion, species can remain stable in spite of frequent hybridization and introgression. While there has been significant progress in understanding this introgression, we still have little overall knowledge of the genomic distribution of introgressed and non-introgressed loci.

6. *Lysandra coridon*, *L. hispana* and *L. albicans*: frequent hybridization everywhere, strong gene flow and yet species remain distinguishable!

For a long time, the chalkhill blue was considered in Europe to be a single species, *L. coridon*. However, in *Polyommatus sensu lato*, species rarely show consistent differences in genitalia or wing pattern. Because of this, complexity in the *coridon* group was recognized initially due to voltinism. In 1916, Verity observed three emergences of *Lysandra* in the hills around Florence, Italy and showed that this was due to the existence of two separate species: one single-brooded, *coridon sensu stricto*, one double-brooded, *hispana* H.-S. Later on, he recognized *L. caelestissima*, univoltine with a distinctive sky-blue colour, from Montes Universales, central Spain. In Spain, the situation is especially confusing: there are single- and double-brooded forms, and bimodal hybrid zones where they overlap. At one time, clear blue hybrids between *L. caelestissima* and *L. albicans* from Montes Universales were also considered a distinct species, *caerulescens*. For a while the number of species recognized varied from one to four; eventually three were recognized on the basis of chromosome number and voltinism (de Lesse, 1960a, 1969):

1. *Lysandra coridon*: widespread, univoltine, with n=88-90, with an isolate in central Spain, *caelestissima*, considered a subspecies with n=87.
2. *L. albicans*, univoltine, Southwestern Spain, n=82.
3. *L. hispana*, central France and Italy to Northern Spain, bivoltine, n=84.

De Lesse (1969) described ssp. *lucentina* (correctly: *semperi* Agenjo, 1968) from the Alicante region, which he referred to *hispana* on the basis of chromosome number (n=84); later it turned out to be univoltine like *albicans*. He also showed that *L. italaglauca*, described as a species from central Italy was actually a rather abundant hybrid between *L. coridon* (n=88) and *L. bellargus* (n=45). This form, of intermediate colour between the greyish of *L. coridon* and the dazzling blue of *L. bellargus*, was identical to *L. x polonus* (Zeller, 1845), formerly mistaken as a good species from Poland and later recognized as a hybrid (Tutt, 1910). These hybrids occur wherever the parent species fly together, although their frequency varies widely. *Lysandra*

coridon is univoltine and flies around August, while *L. bellargus* is bivoltine and flies in May and September; the hybrid flies in late June. The meiosis of these hybrids displays incoherent equatorial plates, strongly suggesting sterility (de Lesse, 1960a). Ironically, a blue species, *Lysandra syriaca*, from the Middle East was for a while mistaken for *polonus* (Lederer, 1858). Tutt (1914), who had earlier deduced that *polonus* was a hybrid, also correctly interpreted *L. syriaca* as a “good” species. By analogy, de Lesse interpreted *L. caeruleascens* as a hybrid, but, in this case, karyotypes are similar and meiosis appears normal. Laboratory hybrids between *L. coridon* and *L. hispana* obtained by Beuret (1957) proved fertile and viable until the F3 generation. Interestingly, individuals from the last generation had the most chromosomes, as in *Antheraea* moths (Nagaraju & Jolly, 1986). Another “blue” hybrid mistaken for a species, famous for the author who described it, *Lysandra cormion* (Nabokov, 1941), turned out to be a *Lysandra coridon* x *Meleageria meleager* hybrid (Smelhaus, 1947, 1948; Schurian, 1991, 1997). Again, hybridization occurs regularly in some regions (Moulinet, Alpes Maritimes, France, Czechoslovakia).

De Bast (1985) followed up de Lesse's work using morphometric analysis on imaginal morphology and wing pattern. He recognized five species, *L. coridon*, *L. caelestissima*, *L. albicans*, *L. hispana* and *L. semperi*. The latter could be referred either to *hispana* via of karyotype and wing pattern or to *albicans* via voltinism; de Bast therefore considered it a separate species. In 1989, Schurian, after breeding experiments, crosses and morphological studies of all instars from egg to imago, recognized only three species, *coridon*, *albicans* and *hispana* (*semperi* was included within *hispana*).

Based on a restricted sample of 15 populations, Mensi *et al.* (1988) separated *coridon* and *caelestissima* as species because of a diagnostic allozyme (*Pk-2-105*), absent in *caelestissima*. Lelièvre (1992) systematically sampled 75 populations, collected by himself and HD, in order to cover all known systematic units and to test for hybrid zones in France and Spain. Allozyme analysis showed that two main entities could be readily distinguished: *coridon* + *caelestissima*, and *hispana* + *albicans* + *semperi*, with Nei's $D \approx 0.05$ between the two groups. In contrast, *L. bellargus* was separated from the *coridon* group by a $D \approx 0.30$. No diagnostic alleles were found between *coridon* and *caelestissima*, contradicting Mensi *et al.*, (1988). Therefore, there is little reason to consider them as separate species. The chief argument for separation is the colour of male imagines, but, in northern Spain, populations are often of intermediate colour (ssp. *manleyi* and *asturiensis*). A sex-limited morph, the blue “*syngrapha*” female, shared by *coridon* and *caelestissima* (Descimon, 1989) also suggests conspecificity. Disjunct distributions of the two taxa prevent use of the cohabitation criterion. A conservative solution is thus to merge all the populations into a single species with some strong subspecies.

The tale of *L. coridon* in Tyrrhenian islands is almost incredible. Its lime-loving foodplant, *Hippocrepis comosa*, is very scarce on the mainly acidic soil of these islands. The description in 1977 of ssp. *nufrellensis* from the remote granitic Corsican Muvrella massif by Schurian attracted scepticism, but was confirmed in 2006 by Schurian *et al.* – Muvrella granite is hyperalkaline and supports *H. comosa*! *L. coridon*, described as *gennargenti*, was also found in Sardinia on more easily accessed calcareous patches (Leigheb, 1987). Both populations are well characterized by adult wing pattern (the males are vivid blue and females are always blue) and preimaginal stages. Marchi *et al.* (1996), using allozyme analysis, left the form as a subspecies of *coridon*. However, Jutzeler *et al.* (2003a, b) did not lose an opportunity

to raise yet another known form to species rank, based only on preimaginal morphology.

In the “*hispana-semperi-albicans*” complex, on the other hand, things are much more complicated. Populations assigned to one of these putative taxa by “classical” criteria (namely, wing pattern, distribution and voltinism) are not distinguishable via allozymes. This is especially true for “*albicans*” and “*semperi*”, which broadly overlap in their allozyme polymorphisms.

Hybrid zones between the taxa give rise to additional complexity. A hybrid zone exists between *caelestissima* and *albicans* in Montes Universales (central Spain); both are single-brooded and fly at the same time of year. The former flies at rather high elevation (1200-1800 m), the latter in lower zones (800-1400 m). They overlap at intermediate altitudes, where putative male hybrids (“*caerulescens*”) can easily be detected by wing colour. We have studied three samples, each containing ~30 individuals: the first from a pure *caelestissima* locality (Paso del Portillo); the second from an *albicans* locality (Carpio del Tajo); and a third area of cohabitation, where hybrid *caerulescens* reach a frequency of 10% or more (Ciudad Encantada). Allozyme genotypes were concordant with colour pattern in 77% of the cohabiting sample. Discordant individuals were all “*caerulescens*”, that is, presumably hybrids, and their allozyme genotypes were intermediate (Lelièvre, 1992). The hybrid zone thus appears more or less bimodal, even though hybrids were rather abundant.

Two other hybrid zones were studied in northern Spain (at Ansó and Atarés in the Jaca region), where single-brooded *L. coridon manleyi* overlaps with double-brooded *L. hispana*. The former species again flies at a higher elevation, but the two overlap at intermediate altitudes. “Pure” reference populations were again studied nearby: Aranqüite and Embalse de Oliana, respectively. In the hybrid zone at Ansó, the variously coloured butterflies were hard to separate genetically. Individuals were either genetically similar to those from one or other pure sample, or intermediates. In the second hybrid zone, at Atarés, two visually different categories of individuals were found, some with the obvious *coridon* phenotype of clear blue colour, the others greyish-white and similar to *hispana*. Intermediate specimens were scarce and none were analysed genetically. Paradoxically, all genotypes from the cohabitation zone, including those classified as *hispana* by wing pattern, corresponded to *coridon* from Aranqüite, rather than to *hispana* from Oliana, so introgression is suspected (Lelièvre, 1992).

More recently, bivoltine *Lysandra* populations flying in Southern Slovakia were separated out as a species, *Polyommatus slovacus* (Vitaz *et al.*, 1997), on the basis of subtle adult morphological differences (the bluish dorsal hue of male wing pattern and slight differentiation of male and female genitalia). A cohabitation criterion was used, since it apparently flies with univoltine *L. coridon* in some localities, although there is no mention of hybrids. There is no known genetic difference between *L. slovacus* and neighbouring populations of *L. coridon* (Schmitt *et al.*, 2005). Voltinism remains the chief character.

In conclusion, there is one rather clear, homogeneous species, *L. coridon*, with strongly differentiated subspecies in Spain (*caelestissima*) and the Tyrrhenian islands (*nufrellensis*); chromosome characters, phenology, as well as allozyme data support the unity of this taxon. The geographically variable male wing colour pattern conforms to this diagnosis, since populations from Northern Spain are intermediate. In contrast, the same criteria do not provide coherent evidence for splitting the *hispana*

complex into several units. The forms *semperi* and *hispana* share the same karyotype (n=84), but the former is univoltine like *albicans*, which, however, has a different chromosome number (n=82). Allozymes have not yet proved very useful. HD has doggedly sought further contact zones between the three taxa of the *hispana* complex, but in vain. Lelièvre's (1992) work was extremely useful, but his premature death prevented a more complete analysis.

7. The *Erebia tyndarus* group: parapatry, hybrid zones and Gause's principle

This group illustrates the use of successively more sophisticated taxonomic criteria, and the difficulties of applying various species concepts; we therefore employ a historical approach. The *tyndarus* group consists of "brassy ringlets" characterized by cryptic grey hind wing undersides, which provide good camouflage in rocky grasslands. Their distribution stretches from Western North America, across the Pacific to Eurasia, and finally to the Asturias in Spain. Until the 20th century, all were considered to belong to a single variable species. In 1898, Chapman piloted the use of male genitalia in *Erebia* and recognized *E. callias* Edwards from North America, and a submontane form from Asia Minor, *E. ottomana* H.-S., as separate species. In 1908, Reverdin studied wing pattern in Western European taxa, and showed that the Alpine forms could be arrayed in two groups, *E. tyndarus* Esper and *E. cassioides* Reiner & Hohenwarth. The latter can also be recognized in the Pyrénées, Apennines, Balkans and Carpathians. He further noted that the southernmost form, *hispania* Butler from the Sierra Nevada, could be grouped with others from the Pyrénées, *goya* Frühstörfer and *rondoui* Oberthür, without elevating them to species rank.

Warren (1936) recognized four species based on male genitalia: *tyndarus*, *cassioides*, *dromulus* Staudinger (from the mountains of Asia Minor) and *callias*, from North America, Central Asia, Elburz and the Caucasus. In 1949, he pointed out that *cassioides* and *rondoui* (previously included with *tyndarus*) overlapped in the Pyrénées and considered this cohabitation evidence for separate species. In 1954, he extended this to *tyndarus s. str.* on the grounds of cohabitation with *cassioides* in the Bernese Alps.

There is a striking feature in the *tyndarus* group: distributions of the taxa are typically parapatric and in a given region, there is only one form. Distributions overlap only in very narrow contact zones. Sometimes, hybrids are found in various proportions (see below); in other cases, hybridization is absent. Mutual exclusion can be attributed to Gause's (1934) principle: "one species per ecological niche". For the BSC, the *tyndarus* group was somewhat distressing: morphological criteria are weak, and ecological differences minimal, as shown by mutual geographic exclusion. Narrow cohabitation with little or no admixture therefore became the main criterion in this group.

Warren never went beyond genitalic characters, but de Lesse and Lorković initiated a synthetic approach using karyotype, morphometrics of genitalia, wing pattern variation, laboratory crosses, and detailed field studies on distribution and hybrid zones. There was great variation in chromosome number: *hispania*, with n=24, stood out from *cassioides* and *tyndarus*, with n=10 throughout their ranges (Lorković, 1949, 1953; de Lesse, 1953). Later, two cryptic species were discovered: *calcaria* Lrk. (n=8), from the Julian Alps, and *nivalis* Lrk. & de Lesse (n=11), limited to upper elevations of the Eastern Alps, where it flies above *cassioides* or *tyndarus* (Lorković, 1949, Lorković & de Lesse, 1954b). In addition, de Lesse (1955a, c) showed that *E. callias* from North America, *E. iranica* and *E. ottomana* from the Middle East,

displayed markedly different karyotypes (n=15, 51, and 40, respectively). De Lesse (1960a) performed morphometric analyses of genitalia. He reinstated wing pattern as a valuable tool if concordant with other characters. In particular, he noticed that the dark hindwing eyespots could be shifted distally in their fulvous surrounds, rather than being centred, enabling one to group the southernmost taxa, *hispania* and *iranica*, also characterized by high chromosome numbers (n=24-25 and 51-52). Recent studies have shown that satyrine eyespot variation often results from important developmental genetic shifts (Brakefield, 2001). Locally adaptive camouflage wing patterns (see above), such as hindwing underside colour, provided less useful criteria.

Lorković (1954) carried out crosses between several taxa (*calcaria* x *cassioides*, *calcaria* x *hispania* and *cassioides* x *ottomana*). All showed genetic and behavioural incompatibility: assortative mating, together with sterility of primary crosses and of F1 hybrids (Lorković & de Lesse, 1954a). However, the taxa used were not the most significant: *ottomana* is notoriously distant from the other members of the group (see below); *calcaria* and *hispania* differ in karyotype (n=8 and 24 respectively) and their ranges are very distant. The most useful test is *calcaria* x *cassioides*: they have identical karyotypes (n=10) and adjacent distributions, but clear incompatibilities were still found.

It was thus important to investigate contact zones and distribution in nature. A complex pattern of allopatric distribution of *hispania* and *cassioides* was found in the Pyrénées (de Lesse, 1953; Descimon, 1957), with very narrow zones of cohabitation. Only a single putative hybrid was captured by Descimon (de Lesse, 1960a) among several hundred individuals in many zones of overlap. In the central Alps, *tyndarus* occurs as an outpost inserted between two disjunct populations of putative “*cassioides*”. In the absence of differences in chromosomes, genitalia and wing pattern provided the only useful criteria. Westwards, in Val Ferret, SW Switzerland and in adjacent Italy, above Courmayeur, populations of *tyndarus* and “*cassioides*” are separated by narrow unoccupied regions (de Lesse, 1952). Near Grindelwald, in the Bernese Oberland, a cohabitation site with phenotypically intermediate individuals was found. On the Eastern end of the *cassioides*-*tyndarus* contact zone, in Niedertahl, Austria, a cohabitation site was found, but hybrids were not found, even though enhanced variability in genitalia suggested introgression (Lorković & de Lesse, 1955).

Erebia nivalis Lrk. & de L., originally considered a smaller high elevation form of *cassioides* (Lorković & de Lesse, 1954b), was raised to species rank after discovery of its peculiar karyotype (n=11). Cohabitation is often observed at the altitudinal boundary between the two, although hybrids are never found. Competitive exclusion is especially convincing: at Hohe Tauern, a different species occurs on each of two isolated massifs (*cassioides* on Weisseck and *nivalis* on Hochgolling); in both cases the entire span of alpine and subalpine zones (1800-2600 m) is occupied, suggesting competitive release (Lorković, 1958). Similarly, in Eastern parts of their distribution, *cassioides* and especially *tyndarus* reach higher elevations in the absence of *nivalis*. The distribution of *nivalis* is broadly fragmented into two parts, in the Austrian Alps, and in a more restricted area in Bernese Oberland. The gap between the two areas occupied by *nivalis* has been colonized by *tyndarus*. In the Grindelwald area, where all three taxa cohabit, *tyndarus* looks like the more aggressive competitor which has eliminated *nivalis* even from high elevation habitats.

A rather clear picture emerges from these studies (Guillaumin & Descimon, 1976): in Europe, the *tyndarus* group includes several well defined species: *ottomana*, *hispania*, *calcaria*, *nivalis*. The *tyndarus*-*cassioides* pair is more puzzling. By now, a

disjunct assemblage of seemingly subspecific forms were recognized as *cassioides*, including populations from the Asturias, the Pyrénées, Auvergne in French Massif Central, Western and Southern Alps, Eastern Alps, the Apennines and some Balkan massifs. The populations referable to *tyndarus* occurred in a continuous distribution inserted like a wedge between *cassioides* populations in the Central Alps. Lorković (1953) proposed that these taxa were examples of an intermediate category, “semispecies” (Lorković, 1953; Lorković & Kiriakoff, 1958). However, in practice, *cassioides* and *tyndarus* were considered separate species by most lepidopterists (e.g. de Lesse, 1960b).

In 1981, Warren published a supplement to his monograph of the genus *Erebia*. Arguing that chromosomes had little systematic value, he relied mainly on male genitalia and arranged the taxa in a somewhat confusing way. This was accentuated because he considered *cassioides* a *nomen nudum*, in spite of the lectotypification of the figure in Reiner and Hohenwarth by de Lesse (1955a) – he considered the figure was inaccurate. He recognized the following European species:

- 1 *tyndarus* – Central Alps.
- 2 *nivalis* – Austrian Alps and Bernese Oberland.
- 3 *aquitania* Frhst. (= *cassioides pro parte*) – Southern Alps, Dolomites, Karawanken, Montenegro, Etruscan Apennines, Mont Blanc range and Pyrénées in part.
- 4 *neleus* Frr. (= *cassioides pro parte*) – Transylvanian Alps, Austria, Rhodope, Macedonia, Central Alps, Pyrénées (part), Roman Apennines, Abruzzi, Auvergne.
- 5 *calcarius* – Julian Alps.
- 6 *hispania* – Sierra Nevada and Pyrénées.
- 7 *ottomana* – considered very distinct from the other members of the group.

The species designated by Warren in the former *cassioides* group lacked zoogeographical coherence compared with those recognized by de Lesse and Lorković. The only serious (partial) support for Warren’s theses was the suggestion that populations of *cassioides s. l.* east of the *tyndarus* wedge could be called *neleus*, and the western ones *aquitania* (von Mentzer, 1960). This prophetic suggestion, making zoogeographic sense, was largely overlooked at the time.

A much firmer position was adopted by Niculescu (1985): an extreme “lumper”, he used only morphological criteria to unite all of the brassy ringlets in a single polytypic species, *tyndarus*. Much earlier, de Lesse (1960a: 57), had warned about the exclusive use of morphology as criteria to delimit species, especially if already known to be labile and if the classification required illogical zoogeographic distributions. However, Gibeaux (1984) claimed he had discovered *E. calcaria* and *E. tyndarus* closely adjacent to *cassioides* in the Col Izoard region of the French Alps, on the base of wing pattern and genitalic morphology, without reference to karyotype, cohabitation and molecular criteria. Lorković (pers. comm. to H.D.) keenly argued that the genitalic characters used by Gibeaux could be explained by individual variation. Wing pattern differences were confined to the strongly selected, taxonomically useless hindwing undersides.

Ten years later, a far more informative study, based on 17 allozyme loci,

largely confirmed the common ground of previous authors: *ottomana*, the *hispania* complex and *nivalis* were very distinct from other members of the group, with Nei's $D > 0.20$ (Lattes *et al.*, 1994). The single available sample of *tyndarus* differed by $D = 0.14$ from the cluster, while “*cassioides*” itself consisted of clearly differentiated “western” and “eastern” *cassioides* groups. Lattes *et al.* attempted to outflank Warren's rejection of the name *cassioides* by designating a neotype; an actual museum specimen from the Austrian Alps – *cassioides s. str.* therefore now refers specifically to the Eastern taxon. Actually, the older valid name for western “*cassioides*” was *arvernensis* Oberthür (type locality: Northern French Massif Central), and we use it instead of *neleus* below. The rather large genetic distance between *hispania s. str.* from Sierra Nevada and *rondoui* and *goya* from the Pyrénées (Nei's $D = 0.16$), added to slight differences in chromosome number ($n=25$ vs. 24 , respectively) led the authors to consider them different species. However, they did not do the same with two *ottomana* samples from the Italian Alps and Southern French Massif Central, even though they were distant by a Nei's D of 0.18 .

Most recently, a study using allozymes and sequence data from two mtDNA genes was carried out on a limited number of populations (Martin *et al.*, 2002; Eastern “*cassioides*,” in particular, was lacking). There were large genetic distances between *ottomana* and *hispania s. l.*, and their monophyly was confirmed; *tyndarus* (3 populations) also proved monophyletic, while *nivalis* formed a strongly supported group together with *calcaria*; divergence at the mtDNA genes averaged 0.34%. The allozyme data showed a similar pattern to that found by Lattes *et al.*: *nivalis* was located at the end of a long branch. In contrast to *tyndarus*, *arvernensis* did not group as a single cluster and appeared paraphyletic. The basal and terminal branches of these trees were well resolved, but the intermediate branches, which should define the phylogenetic relationships between *tyndarus*, *arvernensis*, *nivalis* and *calcaria*, remained unclear. The lack of Eastern *cassioides s. str.* prevented accurate phylogenetic estimation, since we still do not know if this taxon clusters with *arvernensis*, *tyndarus*, or *nivalis* and *calcaria*.

A final and rather ludicrous episode of this tale occurred in the butterfly distribution Atlases for France (Delmas *et al.*, 1999) and Europe (Kudrna, 2002). The former used the correct name *arvernensis* for “western *cassioides*”. The resultant geographic distributions were correctly documented by Kudrna, but this author also reported older literature records from France (as well as from Spain, parts of Switzerland, and Italy) as “*cassioides*”. Hence an extensive but entirely fictitious pseudo-sympatry of the two taxa was reported in the French Alps and Pyrénées, and even in the Northern Massif Central.

8. *Erebia serotina* Descimon & de Lesse, 1953: a hybrid mistaken for a species.

In September 1953, the 19-year-old HD captured two individuals of an unknown *Erebia* at 1000 m elevation in the Pyrenean valley of Cauterets and showed them to H. de Lesse. After careful examination, they concluded that the butterflies belong to an unknown, late-flying species they named *E. serotina* (Descimon & de Lesse, 1953) – a surprising finding in the mid-20th century. Further individuals were captured regularly in the same region over a period of 10 years, always late in the season and at the same elevation (Descimon, 1963). Chromosome study (Descimon & de Lesse, 1954) disclosed a number of $n=18$.

However, the absence of females in a sample of 18 individuals was intriguing;

Bourgogne (1963) suggested that *E. serotina* was a hybrid between *E. epiphron* and *E. pronoe*, both also present in the region and having chromosome numbers of 17 and 19, respectively. This possibility had been rejected by Descimon & de Lesse, since the two species live at a higher elevation than *serotina* (over 1400 m and above the treeline). Moreover, de Lesse and later Lorković (pers. comm. to HD), who examined the histological preparations of *serotina* testes, considered chromosome pairing during meiosis to be normal. The debate was echoed by Riley (1975) and Perceval (1977), with no additional data. Higgins and Riley (1970) included *E. serotina* as “Descimon’s Ringlet” in their field guide, although the species was not mentioned in later editions or other guides.

A few other specimens were captured the same valley (Lalanne-Cassou, 1972, 1989) and 15 km to the West (Louis-Augustin, 1985) and also in the Spanish Pyrénées, always late and at low elevation (Lantero and Jordana, 1981). Warren (1981) was also inclined to the hypothesis of a hybrid, which he considered to be between *epiphron* and *manto*, another Pyrenean species, on the basis of morphology and against the chromosomal evidence – *manto* has $n=29$, which should yield $n=23$ for the hybrid. At this juncture, both “hybrid” and “good species” hypotheses seemed unlikely.

Forty years later, the retired HD again went in pursuit of *serotina* and found several individuals in September 2000 and 2002 close to Bagnères de Luchon, 60 km east of Cauterets (Descimon, 2004). An analysed individual was heterozygous at all diagnostic allozyme loci between *epiphron* and *pronoe*, while mtDNA showed that *epiphron* was the mother (Meglecz *et al.*, in preparation). Therefore, *serotina* is indeed a hybrid between *epiphron* and *pronoe*. Moreover, after a series of hand-pairing crosses, three hybrids similar to wild *serotina* were obtained by Chovet (1998). Bourgogne’s hypothesis was therefore proved correct and the mystery of “Descimon’s Ringlet” solved; the absence of females may be due to arrested growth, while males undergo accelerated development and hatch before the cold season (see the *Papilio* case above). Now, the riddle has moved on towards other questions: why does *serotina* fly at altitudes where its parents do not? Why does it occur regularly in the Pyrénées, but not in other regions of parental contact?

Hybrids are scarce in *Erebia*: apart from the previously mentioned *arvernensis* *x hispania* hybrid, only two other cases have been recorded. The first, *intermedia* Schwms is found in the Grisons, Switzerland; initially mistaken for a variety of *E. epiphron*, it was later shown being a *flavofasciata x epiphron* hybrid (Warren, 1981). The second has been collected only once, from the Carpathians, and was recognized immediately as a *pronoe x medusa* hybrid (Popescu-Gorj, 1974). Taken in late September, like *serotina*, it was similar to it also in its genitalia. In all three cases, at least one of the parents of *serotina*, *E. epiphron* or *pronoe*, is involved.

9. Other cases of “bad” species in European butterflies

Palaeartic butterflies demonstrate many other cases of uncertain or “fuzzy” species (Tolman & Lewington, 1997). Table 1 gives a general overview. These cases suggest some general patterns of “bad” species relations, often involving hybrid zones. Some such zones present ecological frontiers, in particular at boundaries between lowland and montane taxa: *Pieris napi* and *bryoniae*, *Euchloe crameri* and *simplonia*, *Lycaena tityrus* and *subalpina*, *Melitaea parthenoides* and *varia*, *Coenonympha arcania*, *gardetta* and *darwiniana*, *Pyrgus cirsii* and *carlinae*.

Coenonympha darwiniana may actually be a stabilized hybrid between *arcania* and *gardetta*, since it is found at intermediate elevations between the areas where *arcania* and *gardetta* occur (Holloway, 1980, Porter et al., 1995; Wiemers, 1998). In most cases, the limit coincides with the elevation where two broods per year become impossible because of low mean temperature; a similar phenomenon in latitude is found in most areas where *Aricia agestis* meets its congener *artaxerxes*. Very often, there is a gap where neither form is regularly present, perhaps because in this area, a second brood can be triggered by photoperiod, but does not complete its growth before autumn, and fails. Here, a discrete biological response cannot easily track a continuous environmental change. Another striking feature is that differentiation between clearly distinct taxa is often observed in the Alps, while in the Pyrénées similar distribution gaps are observed, but with much weaker genetic differentiation between single- and double-brooded populations (e.g. *L. tityrus* and *M. parthenoides*). The case of *Maculinea alcon* and *M. rebeli* is so complex and the ecology of both taxa has given rise to so many papers that it deserves separate treatment. The case of these blues is the closest in butterflies to “ecological races.” No differences were found at mtDNA or nuclear EF1- α gene sequences (Als *et al.*, 2004). However, we know too little about gene exchange between the populations to locate them with precision on the bad species-good species spectrum (Wynhoff, 1998).

Other repeated patterns in contact zones suggest 'suture zones' (Remington 1957) caused by secondary contact of whole faunas from different Pleistocene or earlier refuges, especially the Iberian ("Atlanto-Mediterranean"), and Italian + Balkans refuges ("Ponto-Mediterranean", de Lattin, 1957). *Iphiclides podalirius* and *feisthameli*, *Pontia edusa* and *daplidice*, *Colias hyale* and *alfacariensis*, *Lycaena alciphron* and *gordius*, *Melitaea athalia* and *celadussa*, and *Melanargia galathea* and *lachesis*, appear to belong to this category. Desert species such as *Papilio saharae* and *Melitaea deserticola* meet with temperate counterparts in Northern Africa, while montane species also provide examples of differentiation in various refuges followed by subsequent contact. A general feature of these contacts is Gausean exclusion and therefore parapatry; the cases of *Erebia pandrose* and *sthenny*, *E. euryale* forms, *pharte* and *aethiopellus* are comparable with the *tyndarus* group in this respect. Finally, Corsican and Sardinian endemics are somewhat different; they might be expected to provide parallels with *P. machaon* and *hospiton*, but they lack genetic differentiation or pre- and post-mating incompatibility; consequently, they are not able to cohabit.

V. General discussion

The examples studied here can serve as a testbed for theories and concepts of species and speciation, and of their use in answering questions such as: Are there one, two, or more “good” species involved, or is this an example of speciation in progress? Can we use the results to suggest a simple and unequivocal, or at least useful nomenclature? Is there a general procedure, using the tools and concepts already mentioned, to allow us to reach this goal?

The simplest case is *E. serotina*. Originally ranked as a species, it ended up as a mere hybrid: 1→0. Here the difficulty was technical: it was finally through the use of molecular markers that the parent species and the sexes involved in the cross were recognized. In the case of *L. polonus* and *L. itala glauca*, the tools were cytological; in these cases, the sex of the parents involved remains unknown, although mtDNA analysis could easily solve the question. Among many other known hybrids (Table 1),

the majority have been identified only *via* wing pattern. There is an opposite case, where a species, *L. syriaca*, was recognised after being initially confused with the hybrid *polonus*: 0→1. Hybridization does not occur in all zones of cohabitation with the same frequency, as seen in all the cases studied here. The behaviour of hybrids can be not only different from either parent, but also not intermediate; this is especially striking with *serotina*, but is also observed with *apollo* x *phoebus* hybrids (Descimon *et al.*, 1989).

With *L. sinapis* and *reali*, we have an opposite, but equally clear case: 1→2. The data provide an unambiguous result under all species concepts: there are clear morphological differences; gene pools are completely isolated (to satisfy BSC adepts); the ecological niches are different and the two species form mutually monophyletic assemblages and thus raise no problem for phylogeneticists.

Things become more complex with *Zerynthia*. Few doubt that *Z. rumina* and *polyxena* are “good” species. Again, there are obvious morphological differences, and there is a rather strong separation of gene pools – hybrids are scarce enough to satisfy BSC groupies, in spite of broad sympatry and character displacement in ecological preferences. Phylogeneticists will be happy that each species constitutes a monophyletic assemblage. However, serious genomic incompatibilities were observed between distant populations within each of these species, especially within *rumina*. In fact, the level of incompatibility between the species was not markedly greater than within each. So does *Zerynthia* contain one, two, three, four, or even more species? These findings occurred only as a result of crosses between forms which do not co-occur naturally; they are artefacts. Similar incompatibility effects have also recently been observed within the well-known tropical species *Heliconius melpomene* (Jiggins *et al.*, 2001). It is wisest to conclude: 2→2.

The situation with *Papilio hospiton* and *P. machaon* is clearer, but fits less easily with theory. Obviously these two constitute “good” species, conforming to morphological, biological and cladistic concepts. *Parnassius apollo* and *phoebus* are a similar case. However, the evidence for some mutual introgression corresponds more closely to the “genic view” of speciation. Meanwhile, the asymmetrical character of introgression in *Papilio* fits less perfectly. It seems likely that these *Papilio* diverged beyond the point of no return in allopatry, and that introgression occurred only after *P. machaon* again became sympatric. The case of *Parnassius apollo* and *P. phoebus* is similar, but the two species seem likely to have been in close proximity for a long time. In this case, gene flow would have been progressively reduced. Yet, in spite of introgression, all four species remain “good,” in the sense of “distinguishably different”, wherever they overlap.

With the brown *Agrodiaetus*, the situation changes. Hybrids are morphologically undetectable. Karyotype becomes questionable, here, as a species criterion, unless one allows the concept of karyospecies (e.g. Wiemers, 2003). Until recently, a karyotype markedly different, either in number or size of chromosomes was taken as proof of species status because chromosomal differences directly provide mixiological incompatibility. On this basis, allopatric populations distinct in chromosome number were separated as “good” species. This assumption has now been questioned by Kandul *et al.* (2004), who observed that sympatric strains with strongly differentiated karyotypes could not be differentiated using molecular markers, and therefore likely belonged to a single species. Frustratingly, karyotype variation in this group is sometimes associated with genetic and phylogenetic

differentiation, and sometimes not. So how many “species” are included in Western taxa of brown *Agrodiaetus*? Clearly, *A. ripartii* and *fabressei*, which occur in sympatry, must be distinct (ironically, they have the same chromosome number, but the karyotypes have different morphology). For the other populations, all allopatric and with very variable chromosome numbers, the question makes little sense. Nonetheless, in his excellent, exhaustive work on *Agrodiaetus* and related genera, Wiemers (2003) firmly comes down on the side of all of the other taxa being separate species.

In *Hipparchia*, it seems clear that the best solution is to ignore the more extreme splitters and adopt a moderate lumpers approach (Cesaroni *et al.*, 1994), but this remains somewhat arbitrary and, again, depends heavily on the status of allopatric units.

The situation observed today in the *Erebia tyndarus* group is typical of the present state of systematics. Taxonomic decisions made during the first half of the twentieth century lacked much biological insight, but the important contribution of genitalic morphology boosted knowledge. After Huxley’s “new systematics”, even those specializing in morphology, like Warren, began to take the biological species concept into account, especially with respect to cohabitation, but also because genitalic differences were assumed to cause mechanical incompatibility during mating. The bulk of progress on the group was, however, made during the 1950s using karyology, in this case a highly efficient tool. Differences between chromosomal morphs are regularly associated with sterility and other deleterious side-effects of hybridization. However, morphometrics, research on contact zones, and laboratory crosses were combined with chromosomal studies in a synthetic approach which continues to elicit admiration. It is worth noting the enormous contribution made by de Lesse and Lorković in this field. Access to most populations required ascending many hundreds of meters on foot. In his synthesis, de Lesse (1960a) provided impressive distribution maps. But while data on the most important contact zones and centres of distribution were published in detail, many distributional data accumulated by de Lesse remained unpublished, and were lost when he died.

Mostly, the polytypic or “biological” species concept was employed. However, a number of pockets of resistance rebelled against any attempt at consensus. The *Erebia tyndarus* and the forms of the *cassioides-arvernensis* complex remain the most contentious. At present, it is clear that the Grindelwald contact forms a “bimodal hybrid zone” (Jiggins *et al.*, 1997). Gene flow might help to explain contradictions between allozyme and mtDNA sequence data elsewhere (Lattes *et al.* 1994; Martin *et al.* 2002). There are large allozyme distances between *nivalis* and the other taxa, and rather slight ones with mtDNA. Indeed, *nivalis* is more of a high elevation species that must experience a markedly different thermal environment. Watt (2003) has demonstrated that “differentiation or uniformity of polymorphic genotype frequencies over space may be driven by strong local selection pressures”; allozyme divergence may not always yield results independent of selection.

What was the contribution of molecular markers to improve species delimitation in *tyndarus* group? Lattes *et al.* (1994) used Nei’s genetic distance to separate *cassioides* from *arvernensis* and *hispania* from *rondoui*, but ignored the larger differences between the two populations of *ottomana*, without any particular justification. The main problem of using genetic distance as a criterion of species is that the threshold level may differ in each group studied (Avice, 1994). Finally attempts to determine the status of allopatric taxa (including experimental crosses) are

rather like division by zero, the cohabitation criterion acting like the denominator that does not exist.

More significant was the much greater utility of molecular data for reconstructing phylogeny distinguishing monophyly from paraphyly. However, a phylogenetic species concept may be difficult to apply in this case. For example, in the tree published in Fig. 4 of Martin *et al.* (2002), *calcaria* and *nivalis* cluster within a group consisting of all the *arvernensis* samples, and together form the sister group to the monophyletic *tyndarus* assemblages. Yet *tyndarus* and *arvernensis* act as separate species, since they meet at a bimodal hybrid zone; this causes a logical anomaly for phylogenetic species, since more basal taxa do not seem to reach species rank, but form a paraphyletic group as far as sexual isolation is concerned (if sexual isolation is considered an apomorphy). Further research will perhaps help to resolve some of the tantalizing questions in this group, but, at present, we must confess an inability to answer precisely the question "how many species are there?" One can propose a spectrum of solutions spanning two extremes: the "lumper's" position, with *ottomana*, *hispania*, *tyndarus*; or the "splitter's" position: with the various, very disjunct strains of *ottomana* as "species", *hispania*, *rondoui*, *arvernensis*, *cassioides*, *tyndarus*, *calcaria*, and *nivalis*. However, the precise decision along this spectrum will always be more or less arbitrary.

Although also complex, the *Lysandra coridon* group case is somewhat clearer. In particular, if the phylogenetic species concept is capable of wreaking havoc on the *Erebia tyndarus* group, Wu's (2001) "genic view of species" aids in understanding puzzling features of the *coridon* group. We have mentioned the low level of allozyme differentiation within and between the species of this group, while habitus and ecological features yield stronger, better supported patterns. One must keep in mind that chromosome number is very high in *Lysandra*. Therefore, each linkage group should be small and, hence, hitch-hiking will affect fewer loci during speciation. A majority of the genome might therefore be exchanged freely, while only regions linked to genes affecting sexual isolation and ecological specialization will be kept distinct by strong selection. Otherwise, in this group, the problem of characterizing species is relatively soluble, provided one cuts some Gordian knots. One example of such a unit is provided by *Lysandra coridon*, which displays a very "open" population structure, with few if any genetic differences even between geographically distant populations (Lelièvre, 1992; Schmitt *et al.*, 2002). The main problems are the isolates at the Southern periphery of its distribution: *caelestissima* in the mountains of central Spain and *nufrellensis-gennargentii* in Corsica and Sardinia. The stumbling block of the absence of cohabitation is again encountered. By far the simplest and most sensible solution based on such data would seem to be to merge all the forms into a single species, *coridon*, with some strong peripheral subspecies. Likewise, the *albicans-hispana-semperi* complex is best considered a single species with some variation in chromosome number (as in *coridon*) and adaptive features such as voltinism, in the absence of a clear indication from hybrid zones. On the contrary, the frequent occurrence of bimodal hybrid zones between populations of the *coridon* unit, as previously defined, and of members of the *albicans* complex precludes merging them into a single "good" – or even "bad" – species unit. This case, in common with the *Erebia tyndarus* group, demonstrates the phenomenon of local mutual exclusion due to similar ecological niches, especially foodplant choice. The criteria of voltinism and chromosome number, ranked highly by de Lesse, proved not much more reliable than other criteria. Therefore, to the question: "how many species", we finally answer

“two only” – a simple answer which unfortunately might fray the tempers of some lepidopterists.

Conclusions

“I have just been comparing definitions of species... It is really laughable to see what different ideas are prominent in various naturalists' minds, when they speak of 'species'. In some resemblance is everything & descent of little weight – in some resemblance seems to go for nothing & Creation the reigning idea – in some descent is the key – in some sterility an unfailing test, with others not worth a farthing. It all comes, I believe, from trying to define the undefinable” (Darwin, 1856). Darwin would have found it even more laughable today: Mayden (1997) enumerated no less than 24 species concepts, most of them recent.

Whether species are material, "real" objects, that exist in the absence of human observers as no other taxonomic rank does, or whether they are only a construction of our mind, is a philosophical problem beyond the scope of this paper. Our aim is to use the totality of the existing evidence to suggest simple, practical solutions to taxonomic problems, and we attempt to avoid further adding to the vast slag-heap of useless concepts and definitions of the indefinable. Darwin used only a loose definition of species but he was an experienced taxonomist, knew a great deal about describing actual species, and it was sufficient to convince his readership of transpecific evolution. We believe that, even today, a pragmatic, taxonomic solution is more productive than attempting to decide whose concept is correct.

Two facts are undeniable:

1. Taxonomic decisions based on biological or polytypic species concepts are still common. For instance, Kandul *et al.* (2004) use the term species to mean reproductively isolated populations. Many groups of organisms considered species are well-behaved and obey not only the BSC, but also *most* definitions of species.
2. However, a significant number of rakish taxa will probably always fail to conform to this species morality. They regularly conduct extramarital affairs and produce illegitimate offspring beyond the boundary of the species.

Rogue taxa such as these are the subject of the present chapter. Perhaps the most surprising conclusion we reach is that, in spite of increasing evidence from these well-known European taxa, in some cases flooding out of multiple laboratories using the most modern techniques, many “bad” species stubbornly remain bad under a variety of species concepts. The existence of such rogues is of course a necessary outcome of gradual Darwinian evolution, and it shouldn't worry us. However, when it comes to placing specimens in drawers or data against a name, bad species are a problem. Unfortunately, constructing a perfect species definition that covers both well-behaved and bad species will almost certainly remain a matter of compromise.

Bernardi (1980) has shown that many a specialist in a given group has tinkered with his own special taxonomic categories to cover this kind of situation. An example is the "semispecies" idea of Lorković and Mayr, but many other examples are

scattered throughout the obscure or forgotten literature. Is the solution to house rogue taxa in a special fuzzy species ghetto? This might have been a good idea if bad species were a homogeneous group; however, as we have seen, the intermediate states are variable. In any case, there is no agreement today about the rank even of the supposedly most objective of taxa, the species itself (Isaac *et al.*, 2004). We thus argue that classical taxonomic ranks – species and subspecies – are all we require, to avoid proliferation of ever more finely divided categories.

Returning to the actual bad species analysed above, let us ignore problem taxa that result from taxonomic error, such as the undetected “good” species *Leptidea reali* or the hybrid *Erebia* “serotina”. In the case of *Zerynthia*, there is intra-species incompatibility, coupled with inter-species compatibility; this was discovered only through artificial crosses of geographically separate populations. Perhaps, therefore, we should proclaim the primacy of observations in natural contact or cohabitation over experimental tests, which can give an inaccurate impression of pre- and post-zygotic compatibility (Mayr, 1963; Mallet, 1995). If geographically and genetically intermediate populations disappear, for some reason, we end up with the problem of allopatric entities (see below). Sometimes divergence is so great that it seems logical to classify allopatric taxa as species. But is it really necessary to consider continental and British strains of *Pararge megera* as different species because they display some genetic incompatibility (Oliver, 1972)? We argue it is more informative not to do so.

In the three papilionids (*Zerynthia*, *Parnassius*, *Papilio*), most people looking at natural populations in zones of overlap would declare each pair of species to be “good,” even when hybridization occurs regularly, but sparsely, in at least some areas of cohabitation. We suggest that the same decision should apply to all other cases of bimodal phenotypic and genotypic distribution where hybrids occur (Jiggins & Mallet, 2000), whether or not actual or potential gene flow (introgression) takes place. Similar decisions may be made without difficulty for parapatric species with a contact zone and limited or exceptional hybridization as in the *Erebia tyndarus* group. In the case of *Lysandra*, *Pontia daplidice* and *edusa*, and probably *Melanargia galathea* and *lachesis*, the presence of a bimodal hybrid zone allows us to consider the taxa in contact as species, but here we are near the boundary condition, because, if hybridization becomes much more frequent, hybrid swarms would result, and overlapping populations would become merged into a single, unimodal population. For *Pontia*, there are divergent opinions: Geiger *et al* (1988) and Wenger *et al.* (1993) consider *daplidice* and *edusa* as (semi-)species, while Porter *et al.* (1997) grant them only subspecies rank.

Allopatric forms separated by major geographic discontinuities give rise to a virtually insoluble difficulty. Here, there is a Gordian knot to cut. Mayr (1942, 1963, 1982) repeatedly justified the BSC as the only “non-arbitrary definition of species,” but even he (1982: 282) admits “the decision whether to call such [allopatric] populations species is somewhat arbitrary”. Sperling (2003) likewise suggested that decisions should be made using information, such as genetic distance or karyotype, from closely related taxa that are in contact. This is essentially already implicit in the argument for the use of “potential” gene flow in the BSC. An absolute threshold of similarity or distance is arbitrary, so no-one should harbour illusions about the “reality” of species delimited by this pragmatic approach. The most important objective is to preserve clarity, parsimony, and stability in nomenclature. Therefore,

endemics on Tyrrhenian or Atlantic islands might often be considered subspecies of mainland species if they are moderately differentiated, and we argue that this solution should be employed as far as possible on parsimony grounds. They should be considered species only if they present clear signs of very strong genetic, morphological and biological differentiation above that expected of related mainland species in contact with close relatives. When it comes to allopatric “karyospecies,” one might wish to follow Wiemers (2003), and give specific rank (especially if strongly divergent at other genetic markers). Even here, use of the same species name with chromosome number placed in parentheses would be as informative; this is typically applied, for example, in *Mus musculus*. In general, decisions about the species status of allopatric neighbours is always somewhat arbitrary, and a lot less interesting than obtaining field or genetic data from hybrid zones and parapatric contact zones, or from unimodal lines. Here, one deals with a concrete phenomenon, rather than an investigation into how many angels fit on the head of a pin?

We therefore argue for revival and a modern, scientific justification of the rather neglected and misused (and perhaps rightly, in many cases, much-maligned) rank of subspecies. Very often, subspecies have been used to describe geographic forms recognizable only to their author, which has led to disrepute. But today there is a refreshing trend among lepidopterists to consider only more strongly distinct forms (in morphology, ecology, or genetics) as subspecies, and to lump more dubious geographic forms as synonyms. These general recommendations provide a useful compromise between description of geographic variation, the needs of modern butterfly taxonomy (for example, see Ehrlich and Murphy, 1984; Sperling, 2003), and Darwin’s pragmatic use of the term species in evolutionary studies.

It is a Sisyphian task to try to give a definitive, irrefutable definition of species, but species will continue to function as useful tools in biology for a long time. To the question raised by the French population geneticist Le Guyader (2002): “must we give up on a species concept?” we answer: “no!” We recommend that researchers of the future study gene exchange in the many hierarchical layers of phenotype, genotype and genome in “bad” species of butterflies. This has been done in only a handful of species, such as the larch bud moth (Emelianov *et al.* 2004). Such studies will be surely much more illuminating about the nature of speciation and evolution at the species level than endless discussions on the “essence” of species.

References

- Adanson M. 1763.** *Familles des plantes*. Vincent, Paris.
- Als TD, Vila R, Kandul NP, Nash DR, Yen SH, Hsu YF, Mignault AA, Boomsma JJ, Pierce, NE. 2004.** The evolution of alternative parasitic life histories in large blue butterflies. *Nature* **432**: 386-390.
- Amiet JL. 2004.** Séparation des niches écologiques chez deux espèces jumelles sympatriques de *Leptidea* (Lepidoptera, Pieridae). *Revue d'Ecologie (Terre et Vie)*, **59**: 433-452.
- Andersson J, Borg-Karlson AK, Wiklund C. 2003.** Antiaphrodisiacs in pierid butterflies: a theme with variation. *Journal of Chemical Ecology* **29**: 1489-1499.
- Andersson M. 1990.** The driving force: species concepts and ecology. *Taxon* **39**: 375-382.
- Arnold ML. 1992.** Natural hybridization as an evolutionary process. *Annual Review of Ecology and Systematics* **23**: 237-261.
- Arnold ML. 1997.** *Natural Hybridization and Evolution*. Oxford: Oxford University Press.
- Arpagaus M. 1987.** Vertebrate insulin induces diapause termination in *Pieris brassicae* pupae. *Roux's Archive for Developmental Biology* **196**: 527-530.
- Asmussen MA, Arnold J, Avise JC. 1989.** The effects of assortative mating and migration on cytonuclear associations in hybrid zones. *Genetics* **122**: 923-934.
- Aubert J, Solignac M. 1990.** Experimental evidence for mitochondrial DNA introgression between *Drosophila* species. *Evolution* **44**: 1272-1282.
- Aubert J, Barascud B, Descimon H, Michel F. 1996a.** Systématique moléculaire des argynnes (Lepidoptera: Nymphalidae). *Comptes Rendus de l'Académie des Sciences de Paris. Série III: Sciences de la Vie* **319**: 647-651.
- Aubert J, Descimon H, Michel F. 1996b.** Population biology and conservation of the Corsican swallowtail butterfly *Papilio hospiton* Gén. *Biological Conservation* **78**: 247-255.
- Aubert J, Barascud B, Descimon H, Michel F. 1997.** Ecology and genetics of interspecific hybridization in the swallowtails, *Papilio hospiton* Gén. and *P. machaon* L., in Corsica (Lepidoptera: Papilionidae). *Biological Journal of the Linnean Society* **60**: 467-492.
- Aubert J, Legal L, Descimon H, Michel F. 1999.** Molecular phylogeny of swallowtail butterflies of the tribe Papilionini (Papilionidae, Lepidoptera). *Molecular Phylogenetics and Evolution* **12**: 156-167.

- Avise JC. 1974.** Systematic value of electrophoretic data. *Systematic Zoology* **23**: 465-481.
- Avise JC. 1994.** *Molecular Markers, Natural History and Evolution*. London: Chapman and Hall.
- Avise JC, Ball RM. 1990.** Principles of genealogical concordance in species concepts and biological taxonomy. In: Futuyma DJ, Antonovics J, eds. *Oxford Surveys in Evolutionary Biology*, Vol. 7. Oxford: Oxford Univ. Press, 45-67.
- Ayala FJ, Powell JR. 1972.** Allozymes as diagnostic characters of sibling species of *Drosophila*. *Proceedings of the National Academy of Sciences, USA* **69**: 1094-1096.
- Balleto E, Toso GG. 1979.** On a new species of *Agrodiaetus* (Lycaenidae) from Southern Italy. *Nota Divulg. Ist. Agr. Univ. Napoli Portici* **2**: 13-22.
- Barton NH, Hewitt GM. 1989.** Adaptation, speciation and hybrid zones. *Nature (London)* **341**: 497-503.
- De Bast, B. 1985.** La notion d'espèce dans le genre *Lysandra* Hemming, 1933 (Lepidoptera, Lycaenidae). *Linneana Belgica* **10**: 98-110, 175-190, 194-208.
- Baum DA, Larson A. 1991.** Adaptation reviewed: a phylogenetic methodology for studying character macroevolution. *Systematic Zoology* **40**: 1-18.
- Beltrán MS, Jiggins CD, Bull V, Linares M, Mallet J, McMillan WO, Bermingham E. 2002.** Phylogenetic discordance at the species boundary: comparative gene genealogies among rapidly radiating *Heliconius* butterflies. *Molecular Biology and Evolution* **19**: 2176-2190.
- Benes J, Konvicka M, Vrabec V, Zamecnik J. 2003.** Do the sibling species of small whites, *Leptidea sinapis* and *L. reali* (Lepidoptera, Pieridae) differ in habitat preferences? *Biologia, Bratislava* **5**: 943-951.
- Berenbaum M. 1995.** Chemistry and oligophagy in the Papilionidae. In: Scriber JM, Tsubaki Y, Lederhouse RC, eds. *Swallowtail Butterflies: their Ecology and Evolutionary Biology*. Gainesville, Florida: Scientific Publishers, 27-38.
- Berlacher SH. 1998.** Origins: a brief history of research on speciation. In: Howard DJ, Berlacher SH, eds. *Endless Forms. Species and Speciation*. New York: Oxford University Press, 3-15.
- Bernardi G. 1980.** Les catégories taxonomiques de la systématique évolutive. *Memoires de la Société Zoologique de France* **40**: 373-425.
- Beuret H. 1957.** Studien über den Formenkreis *Lysandra coridon-hispana-albicans*. Ein Beitrag zum Problem der Artbildung (2 Studie). *Mitt. Ent. Ges. Basel, N.F.* **7**: 17-36, 37-59.
- Bird CD, Hilchie GJ, Kondla NG, Pike EM, Sperling FAH. 1995.** *Alberta*

Butterflies. Edmonton, Alberta: Provincial Museum of Alberta.

Blum MJ. 2002. Rapid movement of a *Heliconius* hybrid zone: evidence for phase III of Wright's shifting balance theory? *Evolution* **56**: 1992-1998.

Bourgogne J. 1953. *Melitaea athalia athalia* Rott. et *M. athalia helvetica* Rühl (*pseudathalia* Rev.) en France. Etude biogéographique (Lep. Nymphalidae). *Annales de la Société Entomologique de France* **72**: 131-176.

Bourgogne J. 1963. Réflexions au sujet d'une espèce singulière, *Erebia serotina* (Nymphalidae: Satyrinae). *Alexanor* **3**: 363-368.

Bowers MD. 1983. The role of iridoid glycosides in host-plant specificity of checkerspot butterflies. *Journal of Chemical Ecology* **9**: 475-493.

Brakefield P. 1990. Case studies in ecological genetics. In: Kudrna O, ed. *Butterflies of Europe*, Vol. 2. Wiesbaden: Aula, 307-331.

Brakefield P. 2001. Structure of a character and the evolution of butterfly eyespot patterns. *Journal of Experimental Zoology* **291**: 93-104.

Brock JP. 1990. Origins and phylogeny of butterflies. In: Kudrna O, ed. *Butterflies of Europe*, Vol. 2. Wiesbaden: Aula, 209-233.

Brower AVZ, Egan MG. 1997. Cladistic analysis of *Heliconius* butterflies and relatives (Nymphalidae: Heliconiiti): a revised phylogenetic position for *Eueides* based on sequences from mtDNA and a nuclear gene. *Proceedings of the Royal Society of London Series B: Biological Sciences* **264**: 969-977.

Brown J. 1976a. Notes regarding two previously described European taxa of the genera *Agrodiaetus* Hübner 1822 and *Polyommatus* Kluck 1801 (Lep.: Lycaenidae). *Entomologist's Gazette* **27**: 77-84.

Brown J. 1976b. On two previously undescribed species of Lycaenidae (Lepidoptera) from Greece. *Entomol. Ber. Amsterdam* **36**: 46-47.

Brown J, Coutsis JG. 1978. Two newly discovered lycaenid butterflies (Lepidoptera: Lycaenidae) from Greece, with notes on allied species. *Entomologist's Gazette* **29**: 201-213.

Bull V, Beltrán M, Jiggins CD, McMillan WO, Bermingham E, Mallet J. 2006. Polyphyly and gene flow between non-sibling *Heliconius* species. *BMC Biology* **4**: 11.

Bush GL. 1969. Sympatric host race formation and speciation in frugivorous flies of the genus *Rhagoletis* (Diptera, Tephritidae). *Evolution* **23**: 237-251.

Carbonell F. 2001. Contribution à la connaissance du genre *Agrodiaetus* Hübner (1822). *A. ahmadi* et *A. khorasanensis*, nouvelles espèces dans le nord de l'Iran. *Linneana Belgica* **18**: 105-110.

- Cesaroni D, Lucarelli M, Allori P, Russo F, Sbordoni V. 1994.** Patterns of evolution and multidimensional systematics in graylings (Lepidoptera: *Hipparchia*). *Biological Journal of the Linnean Society* **52**: 101-119.
- Chapman TA. 1898.** A review of the genus *Erebia* based on an examination of the male appendages. *Transactions of the Entomological Society of London* **3**: 209-240.
- Chovet G. 1998.** *L'accouplement chez les Lépidoptères Rhopalocères: divers aspects éthologiques, morphologiques et physiologiques et leurs implications biosystématiques.* Unpublished Doctorat d'Etat des Sciences Naturelles Thèse, Université Piere et Marie Curie, Paris.
- Cianchi R, Ungaro A, Marini M, Bullini L. 2003.** Differential patterns of hybridization and introgression between the swallowtails *Papilio machaon* and *P. hospiton* from Sardinia and Corsica islands (Lepidoptera, Papilionidae). *Molecular Ecology* **12**: 1461-1471.
- Clarke CA, Larsen TB. 1986.** Speciation problems in the *Papilio machaon* group of butterflies (Lepidoptera, Papilionidae). *Systematic Entomology* **11**: 175-181.
- Clarke CA, Sheppard PM. 1953.** Further observations on hybrid swallowtails. *Supplement to the Entomologist's Record and Journal of Variation* **65**: 1-12.
- Clarke CA, Sheppard PM. 1955.** The breeding in captivity of the hybrid swallowtail *Papilio machaon gorganus* Frühstorfer (F) x *Papilio hospiton* Génè (M). *Entomologist* **88**: 265-270.
- Clarke CA, Sheppard PM. 1956.** A further report on the genetics of the *machaon* group of swallowtail butterflies. *Evolution* **10**: 66-73.
- Clarke CA, Sheppard PM, Willig A. 1970.** The use of ecdysone to break a two and a half year pupal diapause in *Papilio glaucus* female x *Papilio rutulus* male hybrids. *Entomologist* **105**: 137-138.
- Cook LG. 2000.** Extraordinary and extensive karyotype variation: a 48-fold range in chromosome number in the gall-inducing scale insect *Apiomorpha* (Hemiptera: Coccoidea: Eriococcidae). *Genome* **43**: 255-263.
- Coyne JA, Orr HA. 1997.** "Patterns of speciation in *Drosophila*" revisited. *Evolution* **51**: 295-303.
- Coyne JA, Orr HA. 2004.** *Speciation*. Sunderland, Mass.: Sinauer Associates.
- Cracraft J. 1983.** Species concepts and speciation analysis. *Current Ornithology* **1**: 159-187.
- Cracraft J. 1989.** Speciation and its ontology: the empirical consequences of alternative species concepts for understanding patterns and processes of differentiation. In: Otte D, Endler JA, eds. *Speciation and its Consequences*.

Sunderland, Mass.: Sinauer Associates, 28-59.

Cuénot L. 1936. *L'Espèce*. Paris: Doin.

Darwin C. 1856. Letter to J.D. Hooker.

Darwin C. 1859. *On the Origin of Species by Means of Natural Selection, or the Preservation of Favoured Races in the Struggle for Life*. 1st ed. London: John Murray.

Dasmahapatra KK, Blum M, Aiello A, Hackwell S, Davies N, Bermingham EP, Mallet J. 2002. Inferences from a rapidly moving hybrid zone. *Evolution* **56**: 741-753.

Dennis, RLH. 1997. An inflated conservation load for European butterflies: increases in rarity and endemism accompany increases in species richness. *Journal of Insect Conservation* **1**: 43-62.

de Lattin G. 1957. Die Ausbreitungszentren der holarktischen Landtierwelt. *Verhein Deutsche Zoologische Gesellschaft* **1956**: 380-410.

de Lesse H. 1952. Contribution à l'étude du genre *Erebia*. Répartition de *E. tyndarus* Esp. et *E. cassioides* R. et H; dans la partie occidentale du Valais. *L'Entomologiste* **4**: 68-70.

de Lesse H. 1953. Formules chromosomiques nouvelles du genre *Erebia* (Lépid.: Rhopal.) et séparation d'une espèce méconnue. *Comptes Rendus de l'Académie des Sciences de Paris. Série III: Sciences de la Vie* **236**: 630-632.

de Lesse H. 1955a. Nouvelles formules chromosomiques dans le groupe d'*Erebia tyndarus* Esp. (Lépidoptères: Satyrinae). *Comptes Rendus de l'Académie des Sciences de Paris. Série III: Sciences de la Vie* **240**: 347-349.

de Lesse H. 1955b. Distribution holarctique d'un groupe d'espèces du genre *Erebia* (Lépidoptères) récemment séparées d'après leurs formules chromosomiques. *Comptes Rendus de la Société de Biogéographie* **276**: 12-18.

de Lesse H. 1955c. Une nouvelle formule chromosomique dans le groupe d'*Erebia tyndarus* Esp. (Lépidoptères: Satyrinae). *Comptes Rendus de l'Académie des Sciences de Paris. Série III: Sciences de la Vie* **241**: 1505-1507.

de Lesse H. 1956. Une nouvelle formule chromosomique dans le groupe d'*Erebia tyndarus* Esp. (Lépidoptères: Satyrinae). *Comptes Rendus de l'Académie des Sciences de Paris. Série III: Sciences de la Vie* **241**: 1505-1507.

de Lesse H. 1960a. Les nombres de chromosomes dans la classification du groupe d'*Agrodiaetus rippartii* Freyer (Lepidoptera: Lycaenidae). *Revue Française d'Entomologie* **27**: 240-262.

de Lesse H. 1960b. Spéciation et variation chromosomique chez les Lépidoptères

Rhopalocères. *Ann. Sci. Nat. Zoo. Biol. Anim. (Série 12)* **2**: 1-233.

de Lesse H. 1961a. Cohabitation en Espagne d'*Agrodiaetus rippartii* Freyer et *A. fabressei* Oberthür. *Revue Française d'Entomologie* **28**: 50-53.

de Lesse H. 1961b. Les nombres de chromosomes chez *Agrodiaetus dolus* et les espèces voisines. *Alexanor* **2**: 57-63.

de Lesse H. 1966. Variation chromosomique chez *Agrodiaetus dolus* Hübner (Lepidoptera: Lycaenidae). *Annales de la Société Entomologique de France* **5**: 67-136.

de Lesse H. 1969. Le nombre de chromosomes dans le groupe de *Lysandra coridon* Poda. *Annales de la Société Entomologique de France* **5**: 469-522.

Delmas S, Maechler J, Sibert J-M. 1999. *Catalogue permanent de l'entomofaune française. 2. Lepidoptera: Rhopalocera.* Dijon: Union de l'Entomologie Française.

de Puysegur K. 1947. Note sur un accouplement entre *Zerynthia polyxena-creusa* Meig. et *Z. rumina-medesicaste* Ill. *Revue Française de Lépidoptérologie* **11**: 10-15.

de Queiroz K. 1998. The general lineage concept of species, species criteria, and the process of speciation. A conceptual unification and terminological recommendations. In: Howard DJ, Berlocher SH, eds. *Endless Forms. Species and Speciation.* New York: Oxford University Press, 57-75.

de Queiroz K, Donoghue MJ. 1988. Phylogenetic systematics and the species problem. *Cladistics* **4**: 317-338.

Deschamps-Cottin M, Aubert J, Barascud B, Descimon H. 2000. Hybridation et introgression entre "bonnes espèces". Le cas de *Parnassius apollo* et *P. phoebus*. *Comptes Rendus de l'Académie des Sciences de Paris. Série III: Sciences de la Vie* **323**: 287-297.

Descimon H. 1957. Nouvelles données sur la répartition d'*Erebia cassioides* Reiner et Hohenwarth et d'*E. hispania* Butler dans les Pyrénées centrales et occidentales. *Revue Française de Lépidoptérologie* **16**: 48-59.

Descimon H. 1963. Nouvelles captures d'*Erebia serotina*. *Alexanor* **3**: 71-80.

Descimon H. 1980. *Heodes tityrus tityrus* Poda et *H. tityrus subalpina* Speyer (Lycaenidae): un problème de spéciation en milieu alpin. *Nota Lepidopterologica* **2**: 123-125.

Descimon H. 1989. La variation géographique du polymorphisme chez les Lépidoptères: femelles bleues et femelles brunes chez *Lysandra coridon* Poda dans le sud-ouest de l'Europe (Lepidoptera: Lycaenidae). *Alexanor* **16**: 23-41.

Descimon, H., 2002 (2004). *Erebia serotina* : retour aux sources. *Alexanor* **22**: 451-465

- Descimon H, de Lesse H. 1953.** Découverte d'un nouvel *Erebia* dans les Hautes Pyrénées. *Revue Française de Lépidoptérologie* **14**: 119-122.
- Descimon H, de Lesse H. 1954.** Nouvelle note sur *Erebia serotina*. *Revue Française de Lépidoptérologie* **14**: 237-241.
- Descimon H, Geiger H. 1988.** Electrophoretic detection of interspecific hybrids in *Parnassius* (Lepidoptera: Papilionidae). *Génétique, Sélection et Evolution* **20**: 435-440.
- Descimon H, Michel F. 1989.** Expériences d'hybridation intra- et interspécifiques dans le genre *Zerynthia* (Papilionidae). Relativité des critères mixiologiques de l'espèce. *Nota Lepidopterologica* **12(Suppl. 1)**: 28-31.
- Descimon H, Genty F, Vesco J-P. 1989.** L'hybridation naturelle entre *Parnassius apollo* (L.) et *P. phoebus* (F.) dans les alpes du sud (Lepidoptera: Papilionidae). *Annales de la Société Entomologique de France (N.S.)* **25**: 209-234.
- Descimon H, Zimmermann M, Cosson E, Barascud B, Nève G. 2001.** Genetic variation, geographic variation and gene flow in some French butterfly species. *Génétique, Sélection, Evolution* **33 (suppl. 1)** : S223-S249.
- Dethier VG. 1954.** Evolution of feeding preference in phytophagous insects. *Evolution* **8**: 33-54.
- Dobzhansky T. 1937.** *Genetics and the Origin of Species*. New York: Columbia Univ. Press.
- Dodd DMB. 1989.** Reproductive isolation as a consequence of divergence in *Drosophila pseudoobscura*. *Evolution* **43**: 1308-1311.
- Drès M, Mallet J. 2002.** Host races in plant-feeding insects and their importance in sympatric speciation. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* **357**: 471-492.
- Eckweiler W, Häuser CL. 1997.** An illustrated checklist of *Agrodiaetus* Hübner, 1822, a subgenus of *Polyommatus* Latreille, 1804 (Lepidoptera: Lycaenidae). *Nachrichten des entomologischen Vereins Apollo* **16(Suppl.)**: 113-168.
- Ehrlich PR. 1961.** Has the biological species concept outlived its usefulness? *Systematic Zoology* **10**: 167-176.
- Ehrlich PR, Murphy DD. 1983.** Butterflies and biospecies. *Journal of Research on the Lepidoptera* **21**: 219-225.
- Ehrlich PR, Murphy DD. 1984.** On butterfly taxonomy. *Journal of Research on the Lepidoptera* **23**: 19-34.
- Ehrlich PR, Raven PH. 1965.** Butterflies and plants: a study in coevolution.

Evolution **18**: 586-608.

Eisner C. 1966. Parnasiidae-Typen in der Sammlung J.C. Eisner. *Zoologische Verhandelingen* **81**: 190 pp.

Emelianov I, Simpson F, Narang P, Mallet J. 2003. Host choice promotes reproductive isolation between host races of the larch budmoth *Zeiraphera diniana*. *Journal of Evolutionary Biology* **16**: 208-218.

Emelianov I, Marec F, Mallet J. 2004. Genomic evidence for divergence with gene flow in host races of the larch budmoth. *Proceedings of the Royal Society of London Series B: Biological Sciences* **271**: 97-105.

Feder JL. 1998. The apple maggot fly, *Rhagoletis pomonella*: flies in the face of conventional wisdom. In: Howard DJ, Berlocher SH, eds. *Endless Forms. Species and Speciation*. New York: Oxford University Press, 130-144.

Feder JL, Roethele JB, Wlazole B, Berlocher SH. 1997. Selective maintenance of allozyme differences among sympatric host races of the apple maggot fly. *Proceedings of the National Academy of Sciences, USA* **94**: 11417-11421.

Federley H. 1945. Polyploidie und non-disjonction in der Gametogenese einiger Lepidopteran. *Commentationes Biologicae, Societas Scientiarum Fennica* **9**: 1-9.

Feeny P. 1995. Ecological opportunism and chemical constraints on the host associations of swallowtail butterflies. In: Scriber JM, Tsubaki Y, Lederhouse RC, eds. *Swallowtail Butterflies: their Ecology and Evolutionary Biology*. Gainesville, Florida: Scientific Publishers, 9-15.

Forster W. 1956. Bausteine zur Kenntnis der Gattung *Agrodiaetus* Scudd. (Lep.: Lycaen.). *Internationaler Zeitschrift der Wiener Entomologische Gesellschaft* **41**: 42-61, 70-89, 116-212.

Freese, A. & Fiedler, K, 2002. Experimental evidence for specific distinctness of the two wood white butterfly taxa, *Leptidea sinapis* and *L. reali* (Pieridae). *Nota lepidopterologica*, **25**: 39-59.

Friberg M, Bergman M, Kullberg J, Wahlberg N, Wiklund C. 2007. Niche separation in space and time between two sympatric sister species – a case of ecological pleiotropy. *Journal of Animal Ecology* **xxx**: 000-000.

Futuyma DJ. 1986. *Evolutionary Biology*. Sunderland, Mass.: Sinauer.

Futuyma DJ, Keese MC. 1992. Evolution and coevolution of plants and phytophagous arthropods. In: Rosenthal GA, Berenbaum MR, eds. *Herbivores: their Interactions with Secondary Plant Metabolites*, Vol. 2. San Diego: Academic Press, 439-475.

Gamisans J. 1991. *La végétation de la Corse*. Genève: Conservatoire et Jardin Botanique de la Ville de Genève.

- Gause GF. 1934.** *The Struggle for Existence*. Baltimore, Maryland: Williams and Wilkins.
- Geiger H. 1990.** Enzyme electrophoretic methods in studies of systematics and evolutionary biology of butterflies. In: Kudrna O, ed. *Butterflies of Europe*, Vol. 2. Wiesbaden: Aula, 397-436.
- Geiger H, Descimon H, Scholl A. 1988.** Evidence for speciation within nominal *Pontia daplidice* (Linnaeus, 1758) in southern Europe (Lepidoptera: Pieridae). *Nota Lepidopterologica* **11**: 7-20.
- Gibeaux C. 1984.** *Erebia tyndarus* Esper et *E. calcaria* Lorković capturés en France (Lep.: Nymphalidae). *Entomologica Gallica* **1**: 49-60.
- Gilbert, LE. 2003.** Adaptive novelty through introgression in *Heliconius* wing patterns: evidence for a shared genetic "toolbox" from synthetic hybrid zones and a theory of diversification. In: Boggs CL, Watt WB, Ehrlich P, eds. *Butterflies. Ecology and Evolution taking flight*. Chicago: Chicago University Press, 281-318.
- Gillespie JH. 1991.** *The Causes of Molecular Evolution*. Oxford: Oxford Univ. Press.
- Goldschmidt RB. 1932.** Prä oder Postreduktion der Chromosomen? Die Lösung eines alten Problems. *Naturwissenschaften* **20**: 358-362.
- Gómez Bustillo MR, Exposito Hermosa A, Martínez Borrego P. 1979.** Una nueva especie por la ciencia: *Agrodiaetus violetae* (Lep.: Lyc.). *Sociedad Hispano-Luso-Americana de Lepidopterología, Revista de Lepidopterología* **7**: 47-54.
- Gómez Bustillo MR, Fernandez Rubio F. 1974.** Mariposas de la Peninsula Ibérica. Ropaloceros (II). Ministerio de Agricultura. Madrid.
- Grant BR, Grant PR. 1998.** Hybridization and speciation in Darwin's finches. The role of sexual imprinting on a culturally transmitted trait. In: Howard DJ, Berlocher SH, eds. *Endless Forms. Species and Speciation*. New York: Oxford Univ. Press, 404-422.
- Grant PR, Grant BR. 1992.** Hybridization of bird species. *Science* **256**: 193-197.
- Gruha JW, Taylor OR. 1980.** Some characteristics of hybrids derived from the sulfur butterflies, *Colias eurytheme* and *C. philodice*: phenotypic effects of the X-chromosome. *Evolution* **34**: 673-687.
- Guillaumin M, Descimon H. 1976.** La notion d'espèce chez les lépidoptères. In: Bocquet C, Générmont J, Lamotte M, eds. *Les Problèmes de l'Espèce dans le Règne Animal*, Vol. 1. Paris: Société zoologique de France, 129-201.
- Gus R, Schifino MT, Araújo AM. 1983.** Occurrence of localized centromeres in Lepidoptera chromosomes. *Revista Brasileira de Genética* **6**: 769-774.

- Guyot H. 2002.** Découverte d'une nouvelle plante-hôte de *Papilio hospiton* en Corse (Lepidoptera: Papilionidae). *Alexanor* **21**: 285-287.
- Häuser CL, Eckweiler W. 1997.** A catalogue of the species-group taxa in *Agrodiaetus* Hübner, 1822, a subgenus of *Polyommatus* Latreille, 1804 (Lepidoptera: Lycaenidae). *Nachrichten des entomologischen Vereins Apollo* **16(Suppl.)**: 53-112.
- Haldane JBS. 1922.** Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics* **12**: 101-109.
- Harrison RG. 1991.** Molecular changes in speciation. *Annual Review of Ecology and Systematics* **22**: 281-308.
- Harrison RG. 1998.** Linking evolutionary pattern and process. The relevance of species concepts for the study of speciation. In: Howard DJ, Berlocher SH, eds. *Endless Forms. Species and Speciation*. New York: Oxford Univ. Press, 19-31.
- Harrison RG, Bogdanowicz SM. 1997.** Patterns of variation and linkage disequilibrium in a field cricket hybrid zone. *Evolution* **51**: 493-505.
- Hatfield T. 1996.** Genetic divergence in adaptive characters between sympatric species of sticklebacks. *American Naturalist* **149**: 1009-1029.
- Hebert PDN, Cywinska A, Ball SL, deWaard JR. 2003.** Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London Series B: Biological Sciences* **270**: 313-321.
- Held C, Spieth H. 1999.** First experimental evidence of pupal summer diapause in *Pieris brassicae* L.: the evolution of local adaptedness. *Journal of Insect Physiology* **45**: 587-598.
- Hennig W. 1968.** *Elementos de una Sistemática Filogenética (Translation of Grundzüge einer Theorie der phylogenetischen Systematik)*. 2nd ed. (Traducida del alemán por Horstpeter H.G.J. Ulbrich. La revisión técnica estuvo a cargo de Osvaldo Reig). Buenos Aires: Editorial Universitaria de Buenos Aires.
- Hesselbarth G, van Oorschot H, Wagener S. 1995.** *Die Tagfalter der Türkei unter Berücksichtigung der angrenzenden Länder*. Vol. **1-3**. Bocholt: S. Wagener Verlag.
- Hewitt GM. 1988.** Hybrid zones -- natural laboratories for evolutionary studies. *Trends in Ecology and Evolution* **3**: 158-167.
- Higgins LG, Riley ND. 1970.** *A Field Guide to the Butterflies of Britain and Europe*. London: Collins.
- Hillis DM, Moritz C, Mable BK, (eds.). 1996.** *Molecular Systematics*. Sunderland, Mass.: Sinauer Associates.
- Holloway JD. 1980.** *Coenonympha darwiniana* Staudinger (Lepidoptera: Satyridae) – a hybrid origin? *Entomologist's Gazette* **31**: 195-198.

- Isaac NJB, Mallet J, Mace GM. 2004.** Taxonomic inflation: its influence on macroecology and conservation. *Trends in Ecology and Evolution* **19**: 464-469.
- Jiggins CD, Mallet J. 2000.** Bimodal hybrid zones and speciation. *Trends in Ecology and Evolution* **15**: 250-255.
- Jiggins CD, Linares M, Mallet J, Naisbit RE, Salazar C, Yang Z. 2001.** Sex-linked hybrid sterility in a butterfly. *Evolution* **55**: 1631-1638.
- Jordan K. 1896.** On mechanical selection and other problems. *Novitates Zoologicae* **3**: 426-525.
- Jordan K. 1905.** Der Gegensatz zwischen geographischer und nichtgeographischer Variation. *Zeitschrift für Wissenschaftliche Zoologie* **83**: 151-210.
- Jutzeler D, Biermann H, Hesselbarth G, Russo L, Sala G, de Bros E. 1997.** Etudes sur la biologie, la morphologie, et l'éthologie de *Hipparchia sbordonii* Kudrna, 1984, de l'Isola di Ponza (Latium, Italie) et *Hipparchia neapolitana* du Monte Faito (Campanie, Italie) et compléments sur la biologie d'*Hipparchia leighebi* (Kudrna, 1976) (Lepidoptera: Nymphalidae, Satyrinae). *Linneana Belgica* **16**: 105-132.
- Jutzeler D, Casula P, Gascogne-Pees M, Leigheb G, Grill A. 2003a.** Confirmation du status spécifique de *Polyommatus gennargentii* (Leigheb, 1987) de Sardaigne comparé à *Polyommatus coridon* de la région de Schaffhouse (CH) par élevage parallèle (Lepidoptera: Lycaenidae). 2ème partie. *Linneana Belgica* **19**: 149-160.
- Jutzeler D, Leigheb G, Manil L, Villa R, Volpe G. 2003b.** Deux espèces de Lycènes négligées de l'espace tyrrhénien: *Lycaeides bellieri* Oberthür (1910) de Sardaigne et de Corse et *Lycaeides villai* sp. nov. de l'Isle d'Elbe (Lepidoptera: Lycaenidae). *Linneana Belgica* **19**: 65-80.
- Kandul NP, Lukhtanov VA, Dantchenko AV, Coleman JWS, Sekercioglu CH, Haig D, Pierce NE. 2004.** Phylogeny of *Agrodiaetus* Hübner 1822 (Lepidoptera: Lycaenidae) inferred from mtDNA sequences of COI and COII and nuclear sequences of EF1- α : karyotype diversification and species radiation. *Systematic Biology* **53**: 278-298.
- Kimura M. 1968.** Evolutionary rate at the molecular level. *Nature (London)* **217**: 624-626.
- King M. 1993.** *Species Evolution: the Role of Chromosome Change*. Cambridge: Cambridge University Press.
- Klemann P. 1930.** Ein neuer Tagfalterhybrid. *Internationale Entomologische Zeitschrift* **23**: 438-442.
- Kudrna O. 1977.** *A Revision of the Genus Hipparchia Fabricius*. Faringdon: E.W. Classey.

- Kudrna O. 2002.** *The Distribution Atlas of European Butterflies.* (Oedippus, 20: 1-342.) : .
- Lalanne-Cassou B, Lalanne-Cassou C. 1972.** Notes de chasse. *Alexanor* **7**: 240.
- Lalanne-Cassou B, Lalanne-Cassou C. 1989.** Note à propos d'*Erebia serotina* Descimon et de Lesse. *Alexanor* **16**: 52-53.
- Lambert DM, Michaux B, White CS. 1987.** Are species self-defining? *Systematic Zoology* **36**: 196-205.
- Lantero M, Jordana R. 1981.** Primera cita de *Erebia serotina* Descimon & de Lesse. *Boletín de la Asociación Española de Entomología* **4**: 185-189.
- Lattes A, Mensi P, Cassulo L, Balleto E. 1994.** Genotypic variability in Western European members of the *Erebia tyndarus* group (Lepidoptera, Satyridae). *Nota Lepidopterologica* **17**(Suppl. **5**): 93-105.
- Lederer J. 1858.** Noch eine syrische Schmetterlinge. *Wien. Entomol. Monatschr.* **2**: 138.
- Le Guyader H. 2002.** Doit-on abandonner le concept d'espèce? *Courrier de l'environnement de l'INRA* **46**: 51-64.
- Leigheb G. 1987.** *Lysandra coridon* ssp. *gennargenti* nova (Lepidoptera: Lycaenidae), nuovo Licenide della Sardegna. *Bolletino del Museo regionale de Scienze Naturale di Torino* **5**: 447-454.
- Lelièvre T. 1992.** *Phylogénie des Polyommatinae et structure génétique de six espèces du genre Lysandra Hemming (Lepidoptères Lycaenidae).* (Mémoires de la Société Entomologique de France.) Université de Provence: Thèse Doctorat.
- Leraut P. 1990.** Contribution à l'étude des Satyrinae de France. *Entomologica Gallica* **2**: 8-19.
- Lewontin RC. 1974.** *The Genetic Basis of Evolutionary Change.* New York: Columbia Univ. Press.
- Loeliger EA, Karrer F. 2000.** Unusual demonstration of autosomal dominant inheritance of the black coloration of one of America's swallowtails: F2 broods of the hybrid *Papilio polyxenes asterius* with *Papilio machaon gorganus* (Papilionidae). *Nota lepidopterologica* **23**: 40-49.
- Lorković Z. 1941.** Die Chromosomenzahlen in der Spermatogenese der Tagfalter. *Chromosoma* **2**: 155-191.
- Lorković Z. 1949.** Chromosomen-Vervielfachung bei Schmetterlingen und ein neuer Fall fünffacher Zahl. *Revue Suisse de Zoologie* **56**: 243-249.
- Lorković Z. 1953.** Spezifische, semispezifische und rassische Differenzierung bei

Erebia tyndarus Esp. *Travaux de l'Institut de Biologie Experimentale de L'Academie Yougoslave* **11-12**: 163-224.

Lorković Z. 1957. Die Speziationstufen in der *Erebia tyndarus* Gruppe. *Bioloski Glasnik. Periodicum Biologorum* **10**: 61-110.

Lorković Z. 1958. Some peculiarities of spatially and sexually restricted gene exchange in the *Erebia tyndarus* group. *Cold Spring Harbor Symposia on Quantitative Biology* **23**: 319-325.

Lorković Z. 1990. The butterfly chromosomes and their application in systematics and phylogeny. In: Kudrna O, ed. *Butterflies of Europe*, Vol. **2**. Wiesbaden: Aula, 332-396.

Lorković Z. 1994. *Leptidea reali* Reissinger 1989 (= *Lorkovićii* Real 1988), a new European species (Lepid., Pieridae). *Natura Croatica* **2**: 1-26.

Lorković Z, de Lesse H. 1954a. Expériences de croisements dans le genre *Erebia* (Lépidoptères: Satyridae). *Bulletin de la Société Zoologique de France* **79**: 31-39.

Lorković Z, de Lesse H. 1954b. Nouvelles découvertes concernant le degré de parenté d'*Erebia tyndarus* Esp. et *E. cassioides* Hohenw. *Lambillionea* **54**: 58-68, 78-86.

Lorković Z, de Lesse H. 1955. Note supplémentaire sur le groupe d'*Erebia tyndarus* Esp. *Lambillionea* **55**: 55-58.

Lorković Z, Kiriakoff SG. 1958. Proposed insertion in the "Règles" of provisions recognising "semispecies" as a special category for the classification and nomenclature of definite groups of taxa as now proposed to be defined. *Bulletin of Zoological Nomenclature* **15/B (Case 58)**: 1031-1033.

Louis-Augustin J. 1985. Nouvelles captures d'*Erebia serotina* dans les Hautes Pyrénées. *Alexanor* **13**: 348.

Lukhtanov VA, Dantchenko AV. 2002a. Descriptions of new taxa of the genus *Agrodiaetus* Hübner [1822] based on karyotype investigation (Lepidoptera, Lycaenidae). *Atalanta* **33**: 81-107, 224-225.

Lukhtanov VA, Dantchenko AV. 2002b. Principles of the highly ordered arrangement of metaphase I bivalents in spermatocytes of *Agrodiaetus* (Insecta, Lepidoptera). *Chromosome Research* **10**: 5-20.

Lukhtanov VA, Wiemers M, Meusemann K. 2003. Description of a new species of the "brown" *Agrodiaetus* complex from South-East Turkey (Lycaenidae). *Nota Lepidopterologica* **26**: 65-71.

Lukhtanov VA, Kandul NP, Plotkin JB, Dantchenko AV, Haig D, Pierce NE. 2005. Reinforcement of pre-zygotic isolation and karyotype evolution in *Agrodiaetus* butterflies. *Nature* **436**: 385-389.

- Mallet J. 1995.** A species definition for the Modern Synthesis. *Trends in Ecology and Evolution* **10**: 294-299.
- Mallet J. 2001.** Gene flow. In: Woiwood IP, Reynolds DR, Thomas CD, eds. *Insect Movement: Mechanisms and Consequences* (Proceedings of a Symposium at the Royal Entomological Society, London). Wallingford, UK: CAB International, 337-360.
- Mallet J. 2005.** Hybridization as an invasion of the genome. *Trends in Ecology and Evolution* **20**: 229-237.
- Mallet J, Barton N, Lamas G, Santisteban J, Muedas M, Eeley H. 1990.** Estimates of selection and gene flow from measures of cline width and linkage disequilibrium in *Heliconius* hybrid zones. *Genetics* **124**: 921-936.
- Manil L, Diringer Y. 2003.** Excursion entomologique en Corse (juin 2003). Note sur les conséquences des incendies sur une population de *Papilio hospiton* Gén. *Bulletin des Lépidoptéristes Parisiens* **12**: 42-44.
- Manley WBL, Allcard HG. 1970.** *A Field Guide to the Butterflies and Burnet Moths of Spain*. Hampton: E.W. Classey.
- Marchi A, Addis G, Exposito Hermosa V, Crnjar R. 1996.** Genetic divergence and evolution of *Polyommatus coridon gennargenti* (Lepidoptera: Lycaenidae) in Sardinia. *Heredity* **77**: 16-22.
- Martin JF, Gilles A, Lörtscher M, Descimon H. 2002.** Phylogenetics and differentiation of the western taxa of the *Erebia tyndarus* group. *Biological Journal of the Linnean Society* **75**: 319-332.
- Martin J-F, Gilles A, Descimon H. 2003.** Species concepts and sibling species: the case of *Leptidea sinapis* and *Leptidea reali*. In: Boggs CL, Watt WB, Ehrlich P, eds. *Butterflies. Ecology and Evolution Taking Flight*. Chicago: University of Chicago Press, 459-476.
- Mayden RL. 1997.** A hierarchy of species concepts: the denouement in the saga of the species problem. In: Claridge MF, Dawah HA, Wilson MR, eds. *Species: The Units of Biodiversity*. London: Chapman & Hall, 381-424.
- Maynard Smith J. 1966.** Sympatric speciation. *American Naturalist* **100**: 637-650.
- Mayr E. 1942.** *Systematics and Origin of Species*. New York: Columbia Univ. Press.
- Mayr E. 1963.** *Animal Species and Evolution*. Cambridge, Mass.: Harvard University Press.
- Mayr E. 1982.** *The Growth of Biological Thought. Diversity, Evolution, and Inheritance*. Cambridge, Mass.: Belknap.

- Mayr E. 1996.** What is a species, and what is not? *Philosophy of Science* **63**: 261-276.
- Mazel R. 1982.** Exigences trophiques et évolution dans les genres *Euphydryas* et *Melitaea s.l.* (Lep.: Nymphalidae). *Annales de la Société Entomologique de France* **18**: 211-227.
- Mazel R, Leestmans R. 1996.** Relations biogéographiques, écologiques et taxonomiques entre *Leptidea sinapis* Linné et *L. reali* Reissinger en France, Belgique et régions limitrophes (Lepidoptera: Pieridae). *Linneana Belgica* **8**: 317-326.
- Megléc E, Péténian F, Danchin E et al. 2004.** High similarity between flanking regions of different microsatellites detected within each of two species of Lepidoptera: *Parnassius apollo* and *Euphydryas aurinia*. *Molecular Ecology*, **13**, 1693-1700.
- Mensi P, Lattes A, Salvidio S, Balleto E. 1988.** Taxonomy, evolutionary biology and biogeography of South West European *Polyommatus coridon* (Lepidoptera: Lycaenidae). *Zoological Journal of the Linnean Society* **93**: 259-271.
- Mensi P, Lattes A, Cassulo L, Balleto E. 1994.** Biochemical taxonomy and evolutionary relationships in *Polyommatus* (subgenus *Agrodiaetus*) (Lepidoptera: Lycaenidae). *Nota Lepidopterologica* **17(Suppl. 5)**: 105-114.
- Munguira M, Martin J, Perez Valiente M. 1994.** Karyology and distribution as tools in the taxonomy of Iberian *Agrodiaetus* butterflies (Lepidoptera: Lycaenidae). *Nota Lepidopterologica* **17**: 125-140.
- Nabokov V. 1941.** *Lysandra cormion*, a new European butterfly. *Journal of the New York entomological Society* **49**: 265-267.
- Nagaraju J, Jolly MS. 1986.** Interspecific hybrids of *Antheraea roylei* and *A. pernyi*: a cytogenetic reassessment. *Theoretical and Applied Genetics* **72**: 269-273.
- Naisbit RE, Jiggins CD, Mallet J. 2001.** Disruptive sexual selection against hybrids contributes to speciation between *Heliconius cydno* and *H. melpomene*. *Proceedings of the Royal Society of London Series B: Biological Sciences* **268**: 1849-1854.
- Naisbit RE, Jiggins CD, Linares M, Mallet J. 2002.** Hybrid sterility, Haldane's rule, and speciation *Heliconius cydno* and *H. melpomene*. *Genetics* **161**: 1517-1526.
- Nazari V, Zakharov EV, Sperling FAH. 2007.** Phylogeny, historical biogeography and taxonomic ranking of Parnassiinae (Lepidoptera, Papilionidae) based on morphology and seven genes. *Molecular Phylogenetics and Evolution* **42**: 131-156.
- Nei M. 1978.** Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* **89**: 583-590.
- Nève G, Megléc E. 2000.** Microsatellite frequencies in different taxa. *Trends in Ecology and Evolution* **15**: 376-377.

- Nice CC, Shapiro AM. 2001.** Population genetic evidence of restricted gene flow between host races in the butterfly genus *Mitoura* (Lepidoptera: Lycaenidae). *Annals of the Entomological Society of America* **94**: 257-267.
- Niculescu EU. 1985.** Problèmes de morphologie et de taxonomie dans le genre *Erebia* (Lep.: Satyridae). *Bulletin de la Société Entomologique de Mulhouse* **1985**: 1-8.
- Nilsson NH. 1930.** Synthetische Bastardierungsversuche in der Gattung *Salix*. *Lunds Universitets Årsskrift, N.F.* **27**: 1-97.
- Noor MA. 1995.** Speciation driven by natural selection in *Drosophila*. *Nature (London)* **375**: 674-675.
- Noor MAF. 1999.** Reinforcement and other consequences of sympatry. *Heredity* **83**: 503-508.
- Oliver CG. 1972.** Genetic differences between English and French populations of the satyrid butterfly *Pararge aegeria*. *Heredity* **29**: 307-313.
- Oliver CG. 1978.** Experimental hybridization between the nymphalid butterflies *Phyciodes tharos* and *P. campestris montana*. *Evolution* **32**: 594-601.
- Olivier A, van der Poorten D, Puplesiene J, de Prins W. 2000.** *Polyommatus (Agrodiaetus) artvinensis*, stat. nov. and *P. (A.) sigberti*, two vicariant species known so far only from Turkey (Lepidoptera: Lycaenidae). *Phegea* **28**: 57-74.
- Orr HA. 1995.** The population genetics of speciation: the evolution of hybrid incompatibilities. *Genetics* **139**: 1805-1813.
- Pashley DP, Ke LD. 1992.** Sequence evolution in mitochondrial ribosomal and ND1 genes in Lepidoptera: implications for phylogenetic analyses. *Molecular Biology and Evolution* **9**: 1061-1075.
- Paterson HEH. 1985.** The recognition concept of species. In: Vrba ES, ed. *Species and Speciation* (Transvaal Museum Monograph, 4). Pretoria: Transvaal Museum, 21-29.
- Perceval MJ. 1977.** *Erebia serotina* Descimon & de Lesse 1953: a possible hybrid. *Entomological Record and Journal of Variation* **89**: 19-20.
- Perez de Gregorio JJ. 1979.** Nota sobre las especies catalanas del género *Agrodiaetus* Hübner 1823 (Lepidoptera). *Bulletín de la Sociedad Catalana de Lepidopterología* **22**: 7-9.
- Popescu-Gorj A. 1974.** A rare hybrid within the genus *Erebia* (Lep., Satyridae). *Travaux du Musée d'Histoire Naturelle "Gregorie Antipa"* **14**: 267-271.
- Porter AH, Shapiro AM. 1990.** Lock-and-key hypothesis: lack of mechanical

isolation in a butterfly (Lepidoptera: Pieridae) hybrid zone. *Annals of the Entomological Society of America* **83**: 107-114.

Porter AH, Schneider RW, Price BA. 1995. Wing pattern and allozyme relationships in the *Coenonympha arcania* group, emphasising the *C. gardetta-darwiniana* contact area at Bellwald, Switzerland (Lepidoptera, Satyridae). *Nota lepidopterologica* **17**: 155-174.

Porter AH, Wenger R, Geiger H, Scholl A, Shapiro AM. 1997. The *Pontia daplidice-edusa* hybrid zone in northwestern Italy. *Evolution* **51**: 1561-1573.

Poulton EB. 1904. What is a species? *Proceedings of the Entomological Society of London* **1903**: lxxvii-cxvi.

Presgraves DC. 2002. Patterns of postzygotic isolation in Lepidoptera. *Evolution* **56**: 1168-1183.

Réal P. 1962. Un phénomène écologique singulier, mais complexe, l'amphiphénotisme, observé chez les piérides. *Annales Scientifiques de l'Université de Besançon (Série II)* **17**: 87-95.

Réal P. 1988. Lépidoptères nouveaux principalement jurassiens. *Mémoires du Comité de Liaison pour les Recherches Ecofaunistiques dans le Jura* **4**: 1-28.

Reissinger E. 1989. Checkliste Pieridae Duponchel, 1835, der Westpalaearktis (Europa, Nordwestafrika, Kaukasus, Kleinasien). *Atalanta* **20**: 149-185.

Remington CL. 1968. Suture-zones of hybrid interaction between recently joined biotas. *Evolutionary Biology* **1**: 321-428.

Reverdin J. 1908. Variétés et aberrations d'*Erebia tyndarus* dans les Alpes de la Suisse. *Bulletin de la Société Lépidoptérologique de Genève* **1**: 192-215.

Reverdin J. 1920. Note sur *Melitaea athalia* Rott. et diagnose d'une espèce nouvelle. *Bulletin de la Société entomologique de France*, **25**: 319-321.

Reverdin J. 1922. *Melitaea athalia* Rott. et *Melitaea pseudathalia* nova species (?). *Bulletin de la Société lépidoptérologique de Genève* **5**: 24-46.

Rice WR, Hostert EE. 1994. Laboratory experiments on speciation: what have we learned in forty years? *Evolution* **47**: 1637-1653.

Richardson BJ, Baverstock PR, Adams M. 1986. *Allozyme Electrophoresis. A Handbook for Animal Systematics and Population Studies*. London: Academic Press.

Riley ND. 1975. *Erebia serotina* Descimon & de Lesse. *Entomological Record and Journal of Variation* **87**: 266.

Robinson R. 1971. *Lepidoptera Genetics*. Oxford: Pergamon Press.

- Rothschild W, Jordan K. 1906.** A revision of the American Papilios. *Novitates Zoologicae* **13**: 411-752.
- Schluter D. 2000.** *The Ecology of Adaptive Radiation*. New York: Oxford University Press.
- Schluter D. 2001.** Ecology and the origin of species. *Trends in Ecology and Evolution* **16**: 372-380.
- Schmitt T, Seitz A. 2001.** Allozyme variation in *Polyommatus coridon* (Lepidoptera: Lycaenidae) identification of ice-age refugia and reconstruction of post-glacial expansion. *Journal of Biogeography* **28**: 1129-1136.
- Schmitt T, Giessl A, Seitz A. 2002.** Postglacial colonisation of Western Central Europe by *Polyommatus coridon* (Poda, 1761): evidence from population genetics. *Heredity* **88**: 26-34.
- Schmitt T, Varga Z, Seitz A. 2005.** Are *Polyommatus hispana* and *Polyommatus slovacus* bivoltine *Polyommatus coridon* (Lepidoptera: Lycaenidae)? The discriminatory value of genetics in taxonomy. *Organisms, Diversity and Evolution* **5**: 297-307.
- Schurian KG. 1977.** Eine neue Unterart von *Lysandra coridon* Poda (Lep.: Lycaenidae). *Entomologische Zeitschrift* **87**: 13-18.
- Schurian KG. 1989.** Bemerkungen zu "*Lysandra cormion* Nabokov 1941" (Lepidoptera: Lycaenidae). *Nachrichten des entomologischen Vereins Apollo, N.F.* **10**: 183-192.
- Schurian KG. 1991.** Nachtrag zu den "Bemerkungen zu "*Lysandra cormion*" (Lepidoptera: Lycaenidae). *Nachrichten des entomologischen Vereins Apollo, N.F.* **12**: 193-196.
- Schurian KG. 1997.** Freilandexemplare des hybriden *cormion* [= *Polyommatus (Melaegeria) coridon* x *P. (M.) daphnis*] (Lepidoptera: Lycaenidae). *Nachrichten des entomologischen Vereins Apollo, N.F.* **18**: 227-230.
- Schurian KG, Hoffmann P. 1980.** Ein neuer Lycaeniden Hybrid: *Agrodiaetus rippartii* Freyer x *Agrodiaetus menalcas* Freyer (Lep. Lycaenidae). *Nachrichten des entomologischen Vereins Apollo, N.F.* **1**: 21-23.
- Schurian KG, Gascoigne-Pees M, Diringer Y, 2006.** Contribution to the life-cycle, ecology and taxonomy of *Polyommatus (Lysandra) coridon nufrellensis* Schurian (1977) (Lepidoptera: Lycaenidae). *Linneana Belgica* **20**: 180-192.
- Seehausen O. 2003.** Hybridization and adaptive radiation. *Trends in Ecology and Evolution* **19**: 198-207.
- Singer MC, Ng D, Vasco D, Thomas CD. 1992.** Rapidly evolving associations among oviposition preferences fail to constrain evolution of insect diet. *American*

Naturalist **139**: 9-20.

Smelhaus J. 1947. *Polyommatus meleager* ESP. x *coridon* PODA (Lep. Lyc.). *Acta Societatis entomologicae Cechoslovaquiae* **44**: 44-47.

Smelhaus J. 1948. *Polyommatus* hybr. *cormion* Nabokov (*meleager* ESP. x *coridon* PODA) (Lep. Lyc.). *Acta Societatis entomologicae Cechoslovaquiae* **45**: 50-55.

Sneath PHA, Sokal RR. 1973. *Numerical Taxonomy. The Principles and Practice of Numerical Classification.* San Francisco: Freeman.

Sokal RR, Crovello TJ. 1970. The biological species concept: a critical evaluation. *American Naturalist* **104**: 107-123.

Sokal RR, Rohlf FJ. 1981. *Biometry.* 2nd ed. San Francisco: Freeman.

Sperling FAH. 1990. Natural hybrids of *Papilio* (Insecta: Lepidoptera): poor taxonomy or interesting evolutionary problem? *Canadian Journal of Zoology* **68**: 1790-1799.

Sperling FAH. 1993. Mitochondrial DNA variation and Haldane's Rule in the *Papilio glaucus* and *P. troilus* species groups. *Heredity* **71**: 227-233.

Sperling FAH. 2003. Butterfly molecular systematics: from species definitions to higher-level phylogenies. In: Boggs CL, Watt WB, Ehrlich P, eds. *Butterflies. Ecology and Evolution taking Flight.* Chicago: University of Chicago Press, 431-458.

Stebbins GL. 1959. The role of hybridization in evolution. *Proceedings of the American Philosophical Society* **103**: 231-251.

Suomalainen E. 1953. The kinetochore and the bivalent structure in the Lepidoptera. *Hereditas* **39**: 88-96.

Suomalainen E. 1965. On the chromosomes of the geometrid moth genus *Cidaria*. *Chromosoma* **16**: 166-184.

Templeton AR. 1989. The meaning of species and speciation: a genetic perspective. In: Otte D, Endler JA, eds. *Speciation and its Consequences.* Sunderland, Mass., USA: Sinauer Associates, 3-27.

Templeton AR. 1994. The role of molecular genetics in speciation studies. In: Schierwater B, Streit B, Wagner GP, De Salle R, eds. *Molecular Ecology and Evolution: Approaches and Applications.* Basel: Birkhäuser, 455-477.

Templeton AR. 1998. Species and speciation. Geography, population structure, ecology, and gene trees. In: Howard DJ, Berlocher SH, eds. *Endless Forms. Species and Speciation.* New York: Oxford Univ. Press, 32-43.

Thomas CD, Singer MC. 1998. Scale-dependent evolution of specialization in a checkerspot butterfly: from individuals to metapopulations and ecotypes. In: Mopper

S, Strauss S, eds. *Genetic Structure and Local Adaptation in Natural Insect Populations*. New York: Chapman & Hall, 343-374.

Ting C-T, Tsaur S-C, Wu C-I. 2000. The phylogeny of closely related species as revealed by the genealogy of a speciation gene, *Odysseus*. *Proceedings of the National Academy of Sciences, USA* **97**: 5313-5316.

Tolman T, Lewington R. 1997. *Butterflies of Britain and Europe*. (Collins Field Guide.) London: HarperCollins.

Toso GG, Balleto E. 1977. Una nuova specie del *Agrodiaetus* Hübner (Lepidoptera: Lycaenidae). *Annali del Museo Civico di Storia Naturale "Giacomo Doria"* **81**: 124-130.

Troiano G, Balleto E, Toso GG. 1979. The karyotype of *Agrodiaetus humedasa* Toso & Balleto, 1976. *Bollettino della Società Entomologica Italiana* **111**: 141-143.

Tutt JW. 1910. *Agriades polonus* Zeller, mit Bemerkungen über die bekannten Exemplare dieser Form. *Soc. Entomol.* **25**: 3-4.

Tutt JW. 1914. *A Natural History of the British Butterflies, their Worldwide Variation and Geographical Distribution*. Berlin: Stock, London & Friedlander.

Van Valen L. 1976. Ecological species, multispecies, and oaks. *Taxon* **25**: 233-239.

Verity R. 1916. Sur deux *Lycaena* confondus sous le nom de *L. (Agriades) coridon* Poda. *Annales de la Société Entomologique de France* **84**: 504-520.

Vitaz L, Balint Z, Zitnan D. 1997. *Polyommatus slovacus* sp. nov. (Lepidoptera, Lycaenidae): the bivoltine relative of *Polyommatus coridon* in Slovakia. *Entomological Problems* **28**: 1-8.

von Mentzer E. 1960. Über die Spezifität von *Erebia neleus* Frr. und *Erebia aquitania* Frhst. (Lep.: Satyridae). *Entomologisk Tidskrift* **81**: 77-90.

Wahlberg N, Zimmermann M. 2000. Pattern of phylogenetic relationships among members of the tribe Melitaeini (Lepidoptera: Nymphalidae) inferred from mitochondrial DNA sequences. *Cladistics* **16**: 347-363.

Wakeham-Dawson A, Spurdens P. 1994. Anomalous blue butterflies of the genus *Agrodiaetus* Hübner (Lepidoptera: Lycaenidae) in Southern Greece. *Entomologist's Gazette* **45**: 13-20.

Warren BCS. 1936. *Monograph of the genus Erebia Dalman*. London: British Museum (Natural History).

Warren BCS. 1949. Three hitherto unrecognized European species of *Erebia*. *Entomologist* **82**: 97-105.

Warren BCS. 1954. *Erebia tyndarus* Esp. and *Erebia cassioides* Rein. and Hohenw.

(Lep.: Satyridae): two distinct species. *Entomologist's Monthly Magazine* **90**: 129-130.

Warren BCS. 1981. *Supplement to the Monograph of the genus Erebia*. Faringdon, Oxon.: E.W. Classey.

Watt WB. 2003. Mechanistic studies of butterfly adaptations. In: Boggs CL, Watt WB, Ehrlich P, eds. *Butterflies. Ecology and Evolution taking flight*. Chicago: Chicago University Press, 319-352.

Watt WB, Wheat W, Meyer H, Martin JF. 2003. Adaptation at specific loci. VIII. Natural selection, dispersal, and the diversity of molecular-functional variation patterns among butterfly species complexes (*Colias*: Lepidoptera, Pieridae). *Molecular Ecology* **12**: 1265-1275.

Wenger R, Geiger H, Scholl A. 1993. Genetic analysis of a hybrid zone between *P. daplidice* (Linnaeus, 1758) and *P. edusa* (Fabricius, 1777) (Lepidoptera: Pieridae) in Northern Italy. *Proceedings of the Fourth Congress of the European Society for Evolutionary Biology*: 488.

White MJD. 1973. *Animal Cytology and Evolution*. 3rd ed. Cambridge: Cambridge, Univ. Press.

Wiemers M. 1998. Needs reference!!!

Wiemers M. 2003. Chromosome differentiation and the radiation of the butterfly subgenus *Agrodiaetus* (Lepidoptera: Lycaenidae: *Polyommatus*) -- a molecular phylogenetic approach. Ph.D. Thesis, Bonn University. Published at: URL: http://hss.ulb.uni-bonn.de/diss_online/math_nat_fak/2003/wiemers_martin - URN: <http://nbn-resolving.de/urn:nbn:de:hbz:5n-02787>

Wilson R, (ed.). 1999. *Species: New Interdisciplinary Essays*. Cambridge, Mass.: MIT Press.

Wimmers C. 1932. Bastarde unserer europäische Bläulinge. *Entomologische Zeitschrift* **45**: 311-314.

Wynhoff I. 1998. The recent distribution of *Maculinea* species. *Journal of Insect Conservation* **2**: 15-27.

Wu C-I. 2001. The genic view of the process of speciation. *Journal of Evolutionary Biology* **14**: 851-865.

Wynne I, Loxdale H, Brookes C. 1992. Use of a cellulose acetate system for allozyme electrophoresis. In: Berry RJ, Crawford TJ, Hewitt GM, eds. *Genes in Ecology*. Oxford: Blackwell, 494-499.

Zeller P. 1845. *Polyommatus polonus*, eine neue Tagfalterart. *Stettiner Entomologische Zeitung* **3**: 351-354.

Zimmermann M, Aubert J, Descimon H. 1999. Phylogénie moléculaire des Méliées (Lepidoptera: Nymphalidae). *Comptes Rendus de l'Académie des Sciences de Paris. Série III: Sciences de la Vie* **322**: 429-439.

Zimmermann M, Wahlberg N, Descimon H. 2000. A Phylogeny of *Euphydryas* Checkerspot Butterflies (Lepidoptera: Nymphalidae) Based on Mitochondrial DNA Sequence Data. *Annals of the Entomological Society of America* **93** : 347-355.

Appendix. Tools for taxonomic practice at species level in butterflies.

The previous parts of this work presented first the theoretical background of taxonomic work on species, and then a series of analyses of peculiar real cases. To sum up, species are delimited by a series of criteria derived from the concept used and the speciation theory associated with it, with an accent on studies on populations in cohabitation or contact.

There are many different types of datasets that can be used. Wing colour morphology is perhaps the most obvious, and of course in butterflies is extremely important. Ecological, behavioural, and distributional data are also important. Differences in genitalia have often been considered to be significant for reproductive isolation via a “lock-and-key” hypothesis (Jordan, 1896; Porter & Shapiro, 1990). As already seen, genitalic data are useful in certain cases, but not always. Chromosomal data are more often reliable, but they can also be misleading. The same might also be true for pheromonal characters, which can be considered both as organismic and synepigonic, but there is little information on butterflies (but see Andersson *et al.*, 2003).

We here present an overview of currently available methods for gathering and analyzing taxonomic data and conducting biological and statistical studies to establish whether taxa might be species or taxa at some subspecific category. Nomenclatural aspects of species delimitation, however, do not form part of the remit of our chapter.

A. Morphological characters.

Data acquisition. Empirical and intuitive, qualitative observations are still used, but biometrical methods have become more normal. Even with qualitative characters, records of a series of states are often performed. In adults, the hard parts of the exoskeleton are most often studied, and genitalia remain favourite characters since the late 19th Century (Jordan, 1896). Wing pattern variation is used in butterflies predominantly because it is both evolutionarily labile and easy to detect and score, and provides useful data for identification in most cases. A still commonly used method in morphometrics consists of measuring anatomical structures under a microscope with a micrometer (see e.g.: de Lesse, 1960a; Cesaroni *et al.*, 1994). Today, automated measurements employing digital imaging can also be used. Larval characters can also be useful: superficial features (pigmentation, pattern) are commonly used, but chaetotaxy of first instar larvae sometimes provides very significant information. The microstructure of the eggs is a great favourite, especially using scanning electron microscopy (SEM). In using egg sculpturings, one must remember that it is actually an imaginal feature, since it results from the imprint of ovary follicles.

Data analysis. Analysis of morphological data may be performed character by character. It is also possible to integrate a dataset from a sample of individuals in multivariate, or reduced space, analyses (RSA). These methods have been great favourites for the French school of statisticians, long led by Benzecri. Systematists may sometimes be reluctant to use them, but they are powerful when correctly used. It is impossible here even to touch on the principles of these methods; the reader should consult works such as Sneath and Sokal (1973) for details of clustering and ordination methods. In brief, there are three main categories of RSA: principal component analysis, using Euclidean distance, factorial correspondence analysis, using a chi-square-based distance, and factorial discriminant analysis (FDA). The latter seems to be the most appropriate to conduct a study on dataset which may reasonably be

supposed to include two (or more) different species. A frequent criticism of RSA is that these methods are descriptive, rather than inferential statistics. However, with some practice, they are excellent tools for exploring a dataset. Genetic data can also be analyzed in the same way.

B. Chromosome characters

The study of chromosomes in butterflies was for a long time dominated by the work of Lorković (1941) and de Lesse (1960a). Since that time, interest has moved towards other types of genetic markers, but chromosome studies are still useful (e.g. Munguira *et al.*, 1994, Wiemers, 2003). Chromosome counting is typically practised on meiotic cells in the testes during spermatogenesis. Generally rounded, small and numerous, Lepidopteran chromosomes are not gratifying objects of study. In approximately 1000 species of Lepidoptera, the distribution of chromosome numbers proved markedly leptokurtic and asymmetrical, with a strong concentration around the modal number ($n=31$), and an extreme scattering of frequencies for the higher numbers (Robinson, 1971). Some members of *Plebicula* (Lycaenidae) display the highest chromosome numbers in metazoans (190-191 for *P. nivescens*, and for *P. atlantica*), while numbers less than 10 are observed in *Erebia* (de Lesse, 1960a). Supernumerary chromosomes are often seen, especially in Satyridae and HesperIIDae, and may produce pronounced intraspecific variation, in particular in *Plebicula* (de Lesse, 1960a).

The significance of chromosome number variation in butterflies has been widely debated (Lorković, 1941; Robinson, 1971; White, 1973; Kandul *et al.*, 2004). Polyploidy seems unlikely as a general mechanism for chromosome number variation in butterflies, despite Lorković's (1941) views. Centromeric fusion or fission seems a more probable cause of chromosomal number variation (Suomalainen, 1965; White, 1973; King, 1993). This could be due to the structure of the Lepidopteran centromere, which is reportedly "diffuse" (Federley, 1945; Suomalainen, 1953), but see Gus *et al.* (1983). A diffuse centromere may allow some amelioration of damage suffered in chromosomal heterozygotes during meiosis. Another insect group with diffuse centromeres, scale insects, also show large variation in chromosome numbers (Cook, 2000). On the other hand, the modality of chromosome number around 31 throughout the Lepidoptera is not easily accounted for under this scenario (White, 1973). Kandul *et al.* (2004) suggest that instances of enhanced chromosome number variation could result from epidemics of transposable genetic elements.

In practice, chromosome study in butterflies is tedious because spermatogenesis often terminates early in adult life. Even in young males, meiotic metaphase equatorial plates in the spermatids, the most favourable stage for counting, are usually scarce. In addition, chromosomes are usually so highly condensed that little intrachromosomal structure is visible. However, particularly in *Agrodiaetus*, differentiation of larger, so-called macro-chromosomes which vary in number and size has been found useful (de Lesse, 1960b; Munguira *et al.*, 1994, Lukhtanov & Dantchenko, 2002b). Moreover, instead of producing conveniently visible giant polytene chromosomes as in Diptera, Lepidoptera appear to adopt polyploidy as a means of up-regulating gene expression in highly active somatic tissues – far less easy to use as a taxonomic or genetic marker.

Hesselbarth *et al.* (1995) put forward the hypothesis that chromosome fission and fusion could have an influence on adaptive abilities. Species with low chromosome numbers should be associated with greater genome stability and more

supergenic association and therefore adapted to stable environments. Conversely, high chromosome numbers should ease recombination and generate many genotypes promoting adaptation to new or unstable environmental conditions. Wiemers (2003) found absolutely no evidence of such a phenomenon in *Agrodiaetus*, the genus displaying the largest variation in chromosome numbers in butterflies. We suggest another possible effect of high chromosome numbers: by increasing the average rate of recombination, they could limit hitch-hiking of genes causing incompatibility and could therefore ease introgression of “neutral” genes in hybrid belts (e.g. in *Lysandra*).

Karyotypic differences between taxa are often taken as a proof of species-level distinction, and this argument can be legitimate. However, caution must be exercised. Supernumerary, genetically insignificant B-chromosomes are common (de Lesse, 1960a, 1961b), and might sometimes be an indication of hybridization (Wiemers, 1998); moreover, when morphologically and ecologically very similar groups of populations occurring in different areas display different karyotypes, it may be premature to base species separation on chromosomal number, in the absence of other evidence such as molecular studies. The term “chromosome races” (Goldschmidt, 1932) does not seem to have been used explicitly in butterflies, but de Lesse (1966) maintained, within a single species, allopatric populations of *Agrodiaetus dolus* from southern Europe with $N=108, 122$ and 124 ; in contrast, Munguira *et al.* (1995) split the taxa into separate species with different karyotypes. Experiments carried out in moths of the genus *Antheraea* showed that two “species”, *A. roylei* and *pernyi* with $n=18$ and 49 respectively, could be intercrossed for 32 generations with fertility and viability intact (Nagaraju & Jolly, 1986).

C. Molecular characters

The history of molecular systematics can be divided into two major stages: a protein phase and a DNA phase. The former, based mainly on allozyme electrophoresis, became important at the end of the 1960s with studies on *Drosophila* and humans (Avice, 1974; Richardson *et al.*, 1986; Hillis *et al.*, 1996), and played a major role in butterfly systematics from the seventies onwards (Geiger, 1990). The DNA phase really came into its own in the 1990s following the development of the polymerase chain reaction PCR.

Protein data. Since the earliest days, electrophoretic study of protein polymorphism revealed a stunning amount of variation (Lewontin, 1974). A bitter debate on the significance of these observations took place in the sixties and seventies: some championed selection as a cause for polymorphism, while others raised mathematical objections (Kimura, 1968) and argued that it must be neutral. Current experimental (Watt, 2003) and theoretical (Gillespie, 1991) evidence suggests that both selection and neutral evolution may be important; consequently, when using protein variation to study taxonomic units, one must be careful that selected variation affecting ecological parameters, such as foodplants (Feder *et al.* 1997), does not obscure taxonomic conclusions.

Analysis of protein data. A classical method for analysing allozyme data is to reduce the multilocus data by means of a calculation of overall genetic distance (Hillis *et al.*, 1996). This can be used in cluster analyses, and subsequently to phylogenetic inference, but there is no obvious level of genetic distance above which two samples can be confidently considered to be separate species. Nei’s (1978) genetic identity (I) and distance ($D = -\ln I$) is regarded as particularly useful, because it corrects for small

sample size and for multiple “hits”, and so should be proportional to time since divergence under a molecular clock. Closely related species of *Drosophila* may be in the range of Nei’s D of 0.05-0.50 or so (Coyne & Orr, 1997). In European butterflies, the genetic distances between species of the same genus range generally between 0.05 and 0.15 (Aubert *et al.*, 1996b; Geiger 1990; Zimmermann *et al.*, 1999). However, pairs of apparently closely related species may be more distant, and, more surprisingly, other pairs of species may coexist without hybridising, but differ hardly at allozyme loci ($D < 0.01$). Diagnostic loci (fixed for different alleles in each population) are useful for studying hybridisation and gene flow between taxa.

Allozyme studies within species have often attempted to estimate gene flow based on the neutral expectation of gene frequency variation. Firstly, one may estimate the variation of gene frequencies between populations via the use of F_{ST} , the standardized variance of gene frequencies, which measures the fraction of genetic diversity (heterozygosity, H_e) found between populations. If gene frequency variation can be assumed to be a balance between homogenisation via gene flow (m) and local divergence due to genetic drift (proportional to $1/2N_e$, where N_e is the effective population size), then $F_{ST} \approx 1/(1 + 4Nm)$. However, there are many problems with these methods, which allow the estimation only of the combined parameter $N_e m$. They should not be applied in any context other than under equilibrium between genetic drift and gene flow; it does not, for instance, apply in the case of gene flow and hybridization between two species, or between ecologically differentiated taxa (Mallet, 2001), because here selection will be involved in the differentiation (*contra* Porter & Geiger, 1995). In such cases, strong natural selection may lead to rapid equilibration of gene frequencies in the presence of gene flow. A much more useful method is available based on correlations (or linkage disequilibria) between loci diagnostic or with strong frequency differences between hybridising taxa. Hybrid zones, in particular, allow estimation of selection and gene flow separately (Mallet *et al.*, 1990; Porter *et al.*, 1997; Mallet, 2001; Blum, 2002; Dasmahapatra *et al.*, 2002).

A species criterion based on “genotype clusters” (Mallet, 1995) can be viewed as an extension of this multilocus method. Genotypes reach bimodality only when several characters or loci are in tight linkage disequilibrium. One may use “assignment methods”, likelihood or distance-based multivariate statistics (see above under “morphological characters”) to cluster genotypes, to determine whether multilocus gaps between clusters are significant; if so, the clusters can be classified as separate species (Aubert *et al.*, 1997; Feder *et al.*, 1998; Deschamps-Cottin *et al.*, 2000). Newer likelihood or Bayesian methods also allow estimation of the rates of hybridization in a sample of a pair of several, bimodally distributed taxa (Cianchi *et al.*, 2003; Emelianov *et al.*, 2003, 2004).

DNA data. DNA methods have outstripped allozyme electrophoresis, but are still in their infancy compared with what might be possible in a few years. The mitochondrial genome, with a mere 16,000 base pairs has been far the most widely used in butterflies (Pashley and Ke, 1992, Wahlberg and Zimmermann, 2000), and elsewhere. Intraspecific mtDNA sequence polymorphism occurs in certain butterfly species but is absent in other cases. For instance, *Papilio machaon* displays polymorphism throughout its range (Michel, pers. comm.), as do *Euphydryas aurinia* and *Melitaea athalia* (Zimmermann, 2000). In contrast, no variation within *Euphydryas maturna* has been observed across a large range (Zimmermann, *et al.*, 2000). The mitochondrial genome is very sensitive to genetic drift, since it has a N_e four times smaller than that of the nuclear genome. Comparison between closely

related species usually shows 1-2% divergence, but strikingly low differences are observed in some instances: 0.2 % between *Euphydryas maturna* and *E. intermedia*. We therefore do not believe that any particular level of divergence can be used as a suitable benchmark or "DNA barcode" for species status.

Nuclear gene sequences are beginning to be used with some success (e.g. Brower & Egan, 1997; Beltrán et al., 2002), while microsatellite loci have proved disappointingly difficult to obtain in butterflies (Nève and Megléc, 2000; Megléc et al., 2004). Amplified fragment length polymorphisms (AFLPs) can also be used as a very abundant source of "fingerprint" markers in analyses of natural populations, including studies of hybridisation in nature (Emelianov et al., 2004). Nonetheless, while useful in mapping, AFLPs are relatively untried as tools for studying populations.

In summary, marker data, whether morphological, cytological or molecular, have allowed us to search organisms for characters with increasing thoroughness, but are not fundamentally different from one another.

D. Ethological and ecological criteria

Treating ethological and ecological characters together seems hardly justified, since they are heterogeneous. However, they all play an active role both in cohesion *within* species and in maintaining separateness between species. They therefore give access to the very factor, reproductive isolation, important in speciation. We will consider the following most important categories. First, there is the ecological niche and its main constituents: habitat and foodplant choice, phenology and diapause; secondly, sexual behaviour and pheromones; and thirdly, geographic distribution, particularly cohabitation.

According to Gause's principle (1934), if two species occupy the same niche, they will mutually exclude one another and will display parapatric distributions, with very limited cohabitation. These cohabitation zones may not necessarily imply hybridization and/or genetic proximity. Alternatively, if the two species share a large area of sympatry, they must be ecologically differentiated. The main difficulty in using such ecological information is circularity. Very often, field entomologists "feel" that two putative species display subtle differences in habitat use but are unable to develop inferential tests to support their impression. Various parameters of the ecological niches occupied by butterflies frequently crop up in studies of butterfly species:

1) *Larval foodplant choice*. The host plant is perhaps the key niche dimension in the life of a phytophagous insect (Dethier, 1954; Futuyama & Keese, 1992; Feeny, 1995; Berenbaum, 1995), and feeding regime may play a major role in speciation, including in some Lepidoptera (Feder, 1998; Drès & Mallet, 2002). Butterflies are generally oligophagous and change in diet is likely to result in a selective regime that might lead to speciation and adaptive radiation (Ehrlich & Raven, 1965). There is certainly evidence for rapid diet evolution in some taxa, such as the Papilionini (Aubert et al., 1999) or Melitaeini (Mazel, 1982; Singer et al., 1992). These changes may appear spectacular, with switching between plant families common (e.g. Rutaceae to Apiaceae in *Papilio*, Dipsacaceae to Caprifoliaceae and Valerianaceae in *Euphydryas*); however, these unrelated plants almost always have important chemical similarities (Berenbaum, 1995; Bowers, 1983). The evidence for host-related speciation in butterflies is thus somewhat weak (see e.g. Nice & Shapiro, 2001, for a case in the Lycaenidae). In the North American *Euphydryas*, where rapid intraspecific diet

evolution has been observed, new host adaptations normally evolve rapidly in local populations, and drive original preferences and adaptations to extinction, rather than causing speciation (Thomas & Singer, 1998).

2) *Diapause control and voltinism*. A butterfly population is expected to have as many broods as climatic conditions and food availability allow. Intraspecific variation in voltinism is common in species which have a wide range. *Melitaea athalia*, for example, is univoltine in Northern Europe, bivoltine in warm regions with a wet summer, and univoltine again in the Mediterranean and in mountains above 1000 m (see also the *Papilio* paragraph in the present work). On the other hand, sometimes it forms a character presumed to differ at the species level: for example the species *Aricia artaxerxes* is univoltine and occurs in Northern Europe, but is replaced by the bi- or multivoltine species *A. agestis* in Southern Europe. Univoltinism can also be a constitutive character within a taxon: in the genus *Euphydryas*, for example, all species are single-brooded, even under conditions which could allow several broods.

The genetic determination of diapause in Lepidoptera has been studied in few cases, where it apparently involves a number of interacting loci (Held & Spieth, 1999); in other cases, it appears to give a pattern suggesting sex-linked inheritance and few genetic factors. In some cases, crosses between related subspecies or species in the Papilionidae give classic “Haldane’s rule” asymmetry in diapause between males and females, suggesting the importance of Z-linkage (see the *Papilio* section in this chapter and below in this part).

E. Mixiological criteria

“Mixiological” is the term applied, especially in France, to phenotypic and behavioural traits which affect hybridisation and introgression between pairs of taxa. In spite of a heated debate about the use of terms such as “isolating mechanisms” (Lambert *et al.*, 1987; Mallet, 1995), all sides agree that a restriction of gene flow is the key process in speciation in sexual taxa. Since many factors may produce this result, it is normal to aggregate these heterogeneous traits under the same heading, “reproductive isolation” (Mayr, 1963); the two major kinds of reproductive isolation are prezygotic and postzygotic isolation.

1) *Prezygotic barriers* may involve spatial and temporal isolation (habitat choice and phenology), mating behaviour and courtship, pheromone differences, mechanical barriers to pairing, and physiological features of insemination before gametic fusion. Prevailing opinion about their origin is that prezygotic barriers are often formed as a by-product of intraspecific coevolution, with selection maintaining compatibility (as in the “recognition” concept of Paterson, 1985) while the system of mating or reproduction diverges. Another argument is that selection may cause divergence in pre-mating traits as a directly selected process (“reinforcement”) to avoid the production of unfit hybrids. Reinforcement has been much debated (Paterson, 1985, Lambert *et al.*, 1987); however, the phenomenon has been demonstrated in some cases (e.g. Noor, 1995; Lukhtanov *et al.*, 2005), and is suspected in the tropical genus *Heliconius* (Jiggins *et al.*, 2001).

2) *Postzygotic barriers* involve inviability or sterility acting on hybrids from the zygote stage onwards. Hybridization experiments show that hybrids between species are often inviable or sterile. Sterility was demonstrated, for example, using *Drosophila* as a research material (Dobzhansky, 1937), but hybrid sterility has been recognized as early as Buffon’s time (Mayr, 1982). However, hybrid sterility and inviability between taxa considered “good species” is far from general (Darwin, 1859,

and several examples in the present work). Fitness is often reduced in hybrids (Rice & Hostert, 1994), not only in physiology (intrinsic or endogenous selection) but also in ecological adaptations that allow individuals to exploit niches of parental taxa (extrinsic or exogenous selection) (Hatfield, 1996; Jiggins & Mallet, 2000). Crosses in captivity must be considered with utmost caution, since careful rearing and pampering can allow certain experimentally obtained hybrids to survive, while they would undoubtedly die under natural conditions. Conversely, the diseases associated with captivity and promiscuity, or unsuitable breeding conditions, can cause the loss of broods which could have thrived in the wild. This uncertainty allowed wags as Loeliger and Karrer (2000) to cast doubt on earlier results of Clarke and Sheppard (1953, 1955, 1956) and Aubert *et al.* (1997), and to negate the existence of postzygotic incompatibilities between *Papilio machaon* and *P. hospiton* – an extraordinary assertion contradicted by all the evidence!

It has become *de rigueur* to refer to all kinds of hybrid inviability and sterility as Dobzhansky-Muller incompatibilities (Orr, 1995), given that they rarely cause inviability or sterility within species, but only when transferred to another genetic background; in other words, their evil effects result from epistatic incompatibilities between genes. It is likely that hybrid inviability between populations can evolve, paradoxically, without producing fitness problems within populations at any time during its emergence.

Let us finish this work by looking more closely to a striking type of genomic incompatibility we have frequently evoked: Haldane's rule (Haldane, 1922): in hybrids, the heterogametic sex (the one with heterogeneous sex chromosomes, *e.g.* XY) tends to be more sterile or unviable than the homogametic sex (*e.g.* XX). The heterogametic sex is the male in most insects, including *Drosophila*, as well as in mammals. The Lepidoptera and birds are notorious exceptions, having heterogametic females: their sex chromosome formula is ZW in the females and ZZ in males, yet obedience to Haldane's rule in Lepidoptera is as good as or better, in reversed form, as in the species with XX/XY sex-determination (Presgraves, 2002). It is surprising, perhaps, that agreement on the explanation, "dominance theory", of the striking facts of Haldane's rule has been reached only recently: the earliest loci to diverge appear to cause incompatibilities only recessively; thus incompatibilities tend to affect the sex chromosome, and mainly in the heterogametic sex. In agreement with dominance theory, sex-linkage of incompatibilities holds for butterflies, where the female is most strongly affected (Grula & Taylor, 1980; Sperling *et al.*, 1990; Aubert *et al.*, 1997; Jiggins *et al.*, 2001; Naisbit *et al.*, 2002) as well as for *Drosophila*, where it is the male (Coyne & Orr, 1997). It is interesting that the general applicability of Haldane's Rule in the Lepidoptera (Presgraves, 2002) implies that maternally inherited markers, such as mitochondrial DNA or W-chromosomes, will rarely be transmitted between species (Sperling, 1990). Thus, species identification based on mitochondrial "DNA barcodes" may work better for Lepidoptera (Hebert *et al.*, 2003) than in other taxa prone to hybridization and introgression.



Plate I. Bad species of European butterflies 1. [LABELLING SHOWN IN PLATES]

Plate IA: *Polyommatus* s.l. species and their hybrids (*Lysandra* + *Meleageria* + *Plebicula*) (all males).

Top row: *L. coridon*, Cauterets, Hautes Pyrénées, France, 12/08/1964, H. Descimon leg.; "polonus" (*L. coridon* x *L. bellargus*), Quès, Pyrénées-Orientales, France, 15/07/1969, H. Descimon leg.; *L. bellargus*, Cauterets, Hautes Pyrénées, France, 03/09/1965, H. Descimon leg.; *P. dorylas*, Vallecillo, Teruel, Spain, 28/07/1988, H. Descimon leg.

Second row: "cormion" (*L. coridon* x *M. daphnis*), 20/06/1989, bred by Klaus Schurian; "samsoni" (*L. hispana* x *L. bellargus*), Digne, Alpes de Haute Provence, France, 06/08/1923, Dr. Glais leg.; "caerulescens" (*L. caelestissima* x *L. albicans*), Ciudad Encantada, Cuenca, Spain, 08/08/1989, H. Descimon leg.; "caeruleonivescens" (*P. dorylas* x *P. nivescens*), Vallecillo, Teruel, Spain, 28/07/1988, H. Descimon leg.

Third Row: *M. daphnis*, Le Fournel, l'Argentière-la Bessée, Hautes-Alpes, France, 25/07/1966, H. Descimon leg.; *L. hispana* Digne, Alpes de Haute Provence, France, 15/05/1980, H. Descimon leg.; *L. caelestissima*, Ciudad Encantada, Cuenca, Spain, 08/08/1989, H. Descimon leg.; *P. nivescens*, Vallecillo, Teruel, Spain, 28/07/1988, H. Descimon leg.

Plate IB. *Parnassius* species and hybrid (all males).

Top: *P. apollo*, Ravin de Roche Noire, Monêtier les Bains, Hautes Alpes, France, 30/07/1974, H. Descimon leg.; middle: *P. apollo* x *phoebus* hybrid, same locality and collector; bottom: *P. phoebus*. Same locality and collector.

Plate IC. *Anthocaris* species and hybrid (all males).

Top: *A. belia euphenoides*, Auriol, Bouches du Rhône, France, 02/04/1981; middle: *A. belia euphenoides* x *cardamines* hybrid, bred by Z. Lorković, 1938; bottom, *A. cardamines*, Châteauvert, Var, France, 04/04/1981.




Plate II. Bad species of European butterflies 2.

Plate IIA: *Zerynthia* species and hybrid (all males).

Top: *Z. polyxena*, Châteauvert, Var, France, 04/04/1981, H. Descimon leg.; middle: *Z. polyxena* x *Z. rumina* hybrid, bred by H. Descimon, 15/04/82; bottom: *Z. rumina*, Châteauvert, Var, France, 15/04/1980, H. Descimon leg.

Plate IIB: *Erebia* species and hybrid (all males).

Top: *E. epiphron*, Barèges, Hautes-Pyrénées, France, 18/07/1958, H. Descimon leg.; middle: "serotina" (hybrid offspring of *E. pronoe* male x *E. epiphron* female), Pont de la Raillère, Cauterets, Hautes Pyrénées, France, 07/09/1954, H. Descimon leg.; bottom: *E. pronoe* Le Campbasque, Cauterets, Hautes Pyrénées, France, 25/08/1958, H. Descimon leg.

Plate IIC: Distribution of brassy ringlets (*Erebia tyndarus* group) in Europe.

Red = *E. hispania* (*hispania* from Sierra Nevada not distinguished from *rondoui* in the Pyrenees). Sky blue = *E. arvernensis*. Dark blue = *E. cassioides*. Green = *E. tyndarus*. Purple (not very distinctive) = *E. calcaria*. Orange = *E. nivalis*. Dark regions between colour patches = cohabitation zones. Populations from the Italian peninsula are considered to be *E. arvernensis* on the basis of allozymes, while those from the Balkans are assigned to *E. cassioides* somewhat arbitrarily. Crosses represent intersections between latitude and longitude at five degree intervals.

Note 1: The distribution of *hispania* and *cassioides* in the Pyrenees is strongly interdigitated. (The distribution in Spain is not well known; expect more surprises!)

Note 2: *E. nivalis* is fragmented into two nuclei by the distribution of *tyndarus*, which seems more competitive than *cassioides* or *arvernensis*. Competitive exclusion probably explains patchwork distributions and scarcity of overlap.

Note added in proof (12 May 2009):

A paper treating the molecular systematics of the *E. tyndarus* group very thoroughly has just been published (Albre et al. 2008). Based on a large and representative sample, including taxa from across the holarctic, it uses fragments of two mitochondrial genes (*ND5* and *CoII*) to reconstruct phylogeny. It clears up the evolutionary relations of the main holarctic taxa nicely. However, when it comes to what they call the ‘terminal group’ (i.e. the *tyndarus* group s. str.), things become more arguable. The authors recognize that ‘species’ in this group are often paraphyletic assemblages, poorly resolved, in part because of few bases sequenced. However, while they grant species status to *E. nivalis*, *E. calcaria*, *E. tyndarus* and *E. cassioides* they also suggest that *E. macedonica* (Balkans), *E. carmenta* (southern Alps) and *E. arvernensis* constitute ‘good’ species. The authors, after reading an earlier version of our own chapter from the internet, then used it in a manner at variance with its spirit. For instance, we noted that Lattes et al. (1994) do not separate at species level *E. ottomana* from Italy and France. We also noted that the same authors do separate *E. hispania* from Sierra Nevada and Pyrénées, although they are no more genetically distant. Albre et al. seem to use our argumentation in reverse sense, to split even the *ottomana* forms.

This demonstrates that Mayr’s so-called Biological Species Concept (BSC), explicitly referred to by Albre et al., is misleading when used to understand “*species in statu nascendi*”. Old jagged Excalibur does not serve the valiant knights so very well to conquer the Holy Grail of systematics, but instead it drives them to adventures similar to those of Monty Python!

Table 1. Some examples of bad species in European Butterflies, including all verified records of interspecific hybridization in the wild Table 1. Some examples of bad species in European butterflies, including all known records of interspecific hybridization in the wild

Taxonomy

From Tolman and Lewington (1997)

Hybrid frequency

C = Common (~Hardy Weinberg)

F = Frequent >1%

R = Regular <1%

E = Exceptional <0.1%

L = Likely, but no data

Characters enabling detection of hybridization, apart from wing pattern. (- mean does NOT occur)

W- = No wing pattern differences

W+/- = differences not striking enough to allow recognition without especial attention

P = Mate choice differences

D = Diapause

G = Genitalia

C = Chromosomes

A = Allozymes

M = Molecular (nuclear & mitochondrial DNA)

E = Ecological

H = Haldane's rule

H⁻¹ = Inverse Haldane's rule

H⁰ = Non-Haldane rule inviability

I = Inviability (e.g. H⁻¹(I))

S = Sterility

Species 1	Species 2	Location	Hybrid frequency	Characters, except morphology	Taxonomic interpretation	Source
<i>Papilio machaon</i>	<i>P. hospiton</i>	Corsica, Sardinia	F	D, A, M, E, H(I), P-	Sibling species	See text
<i>Iphiclides p. podalirius</i>	<i>I. p. feisthamelii</i>	Languedoc	F	A, M	Parapatric sibling species	See text
<i>Zerynthia polyxena</i>	<i>Z. rumina</i>	Provence	R/E	P, G, C, A, M, E, H! I! S!	Parapatric species	See text
<i>Parnassius apollo</i>	<i>P. phoebus</i>	Throughout the Alps	F	G, A, M, H(S)	Partially sympatric species	See text
<i>Artogeia napi</i>	<i>A. bryoniae</i>	Alps	C/F	C, E, H ⁻¹ (I)	Parapatric sibling species	Bowden, 1966; Geiger and Shapiro, 1992; Porter and Geiger, 1995
<i>Artogeia napi</i>	<i>A. balcana</i>	Balkans	?		?Parapatric subspecies	Tolman and Lewington, 1997
<i>Artogeia napi</i>	<i>A. rapae</i>	Britain, Germany	E	A	Sympatric species	Klemann, 1930; Heslop Harrison, 1951
<i>Pontia daplidice</i>	<i>P. edusa</i>	Coastal S. France, Italy	F	A, H ⁰ (I), G	Parapatric sibling species (narrow overlap)	Geiger et al., 1988 ["semispecies"]; Wenger et al., 1993 ["semispecies"]; Porter et al., 1997 [regard as subspecies]
<i>Euchloe crameri</i>	<i>E. simplonia</i>	Alpes Maritimes, Pyrénées	E to F? Not studied!	W+/-A, E	Parapatric species, ecologically divergent forms (montane vs. lowland)	Lux, 1987
<i>Anthocaris belia euphenoides</i>	<i>A. cardamines</i>	S. France, Spain	R/E	H	Partially sympatric species	R. Legras, pers. comm. (Guillaumin and Descimon, 1976)
<i>Anthocaris damone</i>	<i>A. gruneri</i>	Greece	E	A	Partly sympatric species	Rougeot, 1977
<i>Colias crocea</i>	<i>C. erate</i>	Greece, C. Europe	R/F		Partially sympatric species	Röber, 1909; Alberti, 1943; Gere, 1995
<i>Colias hyale</i>	<i>C. erate</i>	C. Europe	R	A	Partially sympatric species	Röber, 1909; Alberti, 1943; Gere, 1995
<i>Colias crocea</i>	<i>C. hyale</i>	Only in lab But likely to occur	E	Confused with aberrant <i>crocea</i> ?	Sympatric species	Ryszka, 1949; Descimon unpubl. (Guillaumin and Descimon, 1976)

<i>Colias myrmidone</i>	<i>C. hyale</i>	E-Europe	E	A, M	Partially sympatric species	Mecke, 1923; Guillaumin and Descimon, 1976
<i>Colias crocea</i>	<i>C. phicomone</i>	Alps	R	E	Partially sympatric species	Descimon, unpub. (Guillaumin and Descimon, 1976)
<i>Colias palaeno</i>	<i>C. phicomone</i>	Alps	E	E	Partially sympatric species	G. Poluzzi, pers. comm. (Guillaumin and Descimon, 1976)
<i>Colias hecla</i>	<i>C. tyche</i> (= <i>nastes</i>)	Norway, Sweden	F/R	"christiernsonni" Lampa	Sympatric species	Kaisila, 1950
<i>Gonepteryx rhamni</i>	<i>G. cleopatra</i>	S. Europe	E	E	Partially sympatric species	Descimon, unpub. (Guillaumin and Descimon, 1976)
<i>Leptidea sinapis</i>	<i>L. reali</i>	Europe	None known	W-, P, G, A, M, E	Partially sympatric species	no evidence for hybridization
<i>Lycæna tityrus subalpina</i>	<i>Lycæna hippothoe</i>	French Alps	R	E!	Partially sympatric species	G. Bernardi, pers. comm. to HD (Guillaumin and Descimon, 1976),
<i>Lycæna tityrus tityrus</i>	<i>Lycæna t. subalpina</i>		C	E, H(I)	Parapatric strong ssp.	Higgins and Riley, 1970; Descimon, 1980
<i>Cupido minimus</i>	<i>E. alcetas</i>	W France	E		Sympatric species	d'Aldin, 1933
<i>Aricia agestis</i>	<i>A. artaxerxes</i>	UK possible elsewhere	E (ancient)	A, M, D, H(I)	Narrowly overlapping sympatric species	I. Wynne & Mallet, unpub.
<i>Agrodiaetus damon</i>	<i>A. ripartii</i>	Balkans	R	A, M	Partially sympatric species	Schurian and Hoffmann, 1975
<i>Agrodiaetus damon</i>	<i>Meleageria daphnis</i>	Alps	E	C, A, M, E "meledamon"	Intergeneric hybrid	Rebel, 1920
<i>Agrodiaetus damon</i>	<i>Polyommatus icarus</i>	Alps	E	A, M "bion"	Intergeneric hybrid	Rebel, 1929a
<i>Lysandra coridon</i>	<i>L. bellargus</i>	Europe	F	C, G, D, A "polonus" Zeller	Sympatric spp.	See text
<i>Lysandra hispana</i>	<i>L. bellargus</i>	S. France, Spain, Italy	R (rarer than polonus)	C, A, M "samsoni"	Sympatric sp.	Cameron-Curry et al., 1980
<i>Lysandra bellargus</i>	<i>L. albicans</i>	SW Spain	R	D, C, A, M	Distant species	Gómez Bustillo and Fernández Rubio, 1974
<i>Lysandra coridon coelestissima</i>	<i>L. albicans</i>	Central Spain	F	C, A, M "caerulescens" Vty.	Partly sympatric species	See text
<i>Lysandra coridon</i>	<i>Agrodiaetus damon</i>	Alps	E	G, C, A, M "corydamon"	Distant species	Rebel, 1929b; Descimon, unpub.

<i>Lysandra coridon</i>	<i>Meleageria daphnis</i>	Alps	R	C, A, M "cornion" Nabokov	Distant species	See text
<i>Lysandra albicans</i>	<i>Plebicula escheri</i>	Spain	E	C, A, M, E	Intergeneric	Decarpentrie, 1977
<i>Lysandra coridon</i>	<i>Polyommatus icarus</i>	Germany	E	C, A, M, E	Intergeneric	Herrmann, 1926
<i>Lysandra coridon</i>	<i>Plebicula dorylas</i>	France	E	C, A, M, E	Intergeneric	Goodman et al., 1925
<i>Plebicula dorylas</i>	<i>Plebicula nivescens</i>	Central Spain	R	C, A, M "caeruleo- nivescens" Verity in G-B and F-R 1974	Partly sympatric related species	Gómez Bustillo and Fernández Rubio, 1974; Descimon, unpub. (Guillaumin and Descimon, 1976)
<i>Polyommatus icarus</i>	<i>P. eros</i>	Alps	E	A, M	Related species, sympatric on mountains	Descimon, unpub.
<i>Polyommatus icarus</i>	<i>Plebejus argus</i>	Germany	E	A, M, E	Intergeneric hybrid	Peter, 1928
<i>Maculinea alcon</i>	<i>M. rebeli</i>	All Europe	?	M, E	Good species or ecological races?	Wynhoff, 1998
<i>Boloria pales</i>	<i>B. napaea</i>	French Alps	F/R?	W-, G, A, M	Partially sympatric species	Descimon, unpub.
<i>Euphydryas aurinia</i>	<i>E. desfontainii</i>	Spain	R	G, A, M, E, HS	Partially sympatric species	de Lajonquière, 1966
<i>Mellicta athalia athalia</i>	<i>M. athalia celadussa</i>	Central France [?]	C	G, W~	Parapatric subspecies?	See text
<i>Mellicta athalia</i>	<i>M. deione</i>	Provence	E	G, A, M	Partly sympatric species	Descimon, unpub.
<i>Mellicta parthenoides</i>	<i>M. varia</i>	Southern FrenchAlps...	F/R	C, H(I)-	Parapatric sibling species	Bernardi, pers. comm. (Guillaumin and Descimon, 1976)
<i>Melanargia russiae</i>	<i>M. lachesis</i>	Eastern Pyrenees	E	A	Species	Tavoillot, 1967
<i>Melanargia galathea</i>	<i>M. lachesis</i>	France and Spain	F (in some overlaps only)		Parapatric sibling species	Higgins, 1969; Wagener, 1984; Essayan, 1990
<i>Hipparchia semele</i>	<i>H. (senthes?) balletoi</i>	Italy	R	G, A	Parapatric sibling species	Sbordoni, pers. comm. But see text
<i>Erebia flavofasciata</i>	<i>E. epiphron</i>	Alps	R	G	Partially sympatric species	See text
<i>Erebia pharte</i>	<i>E. epiphron</i>	Alps	E	G	Sympatric species	Descimon, unpub. (Guillaumin and Descimon, 1976)

<i>Erebia pronoe</i>	<i>E. epiphron</i>	Pyrénées	R	H(1) "serotina" Descimon & de Lesse	Sympatric species	See text
<i>Erebia pronoe</i>	<i>E. medusa</i>	Carpathians	E		Distant species	See text
<i>Erebia cassioides</i>	<i>E. hispania</i>	Pyrénées	R, several zones in the Pyrenees	G, C, A, M	Parapatric sibling species	See text
<i>Erebia cassioides</i>	<i>E. tyndarus</i>	Alps	F	A, M	Parapatric sibling species	See text
<i>Erebia cassioides</i>	<i>E. nivalis</i>	Alps	F	A, M, E	Parapatric sibling species	See text
<i>Coenonympha arcania</i>	<i>C. hero</i>	Northern Europe	F (<i>hero</i> nearly extinct)		Partly sympatric species	Plantrou, 1951; Groß, 1957; R. Legras, pers. comm. (Guillaumin and Descimon, 1976)
<i>Coenonympha darwiniana</i>	<i>C. gardetta</i>	Alps	F	A <i>C. darwiniana</i> itself may be stabilised hybrid <i>gardetta</i> x <i>arcania</i>	Parapatric species	See text

Table References

- Alberti B. 1943.** Zur Frage der Hybridisation zwischen *Colias erate* Esp., *hyale* L. und *edusa* F. und über die Umgrenzung der 3 Arten. *Mitteilungen der Münchner Entomologischen Gesellschaft* **33**: 606-625.
- Bowden SR. 1966.** 'Sex-ratio' in *Pieris* hybrids. *Journal of the Lepidopterists' Society* **20**: 189-196.
- Cameron-Curry V, Leigh G, Cameron-Curry P. 1980.** Due ibridi di *Lysandra bellargus* Rott. (Lepidoptera, Lycaenidae). *Bollettino della Società Entomologica Italiana* **112**: 41-42.
- d'Aldin A. 1933.** Un hybride de *Lycaena* en Dordogne. *Procès-Verbaux. Actes de la Société Linnéenne de Bordeaux* **85**: 142-143.

- de Lajonquière Y. 1966.** À propos de l'*Euphydryas desfontainii* Godart et de ses sous-espèces. *Alexanor* **4**: 347-353.
- Decarpentrie M. 1977.** Communication [sur l'accouplement de *Lysandra albicans* m. x *Plebicula escheri* f.]. *Bulletin du Cercle des Lépidopteristes de Belgique* **6**: 86-87.
- Descimon H. 1980.** *Heodes tityrus tityrus* Poda et *H. tityrus subalpina* Speyer (Lycaenidae): un problème de spéciation en milieu alpin. *Nota Lepidopterologica* **2**: 123-125.
- Essayan R. 1990.** Contribution lépidoptérique à la cartographie des invertébrés européens. XVII. La cartographie des satyrines de France (*Erebia* non compris) (Lep., Nymphalidae, Satyrinae). *Alexanor* **16**: 291-328.
- Geiger H, Descimon H, Scholl A. 1988.** Evidence for speciation within nominal *Pontia daplidice* (Linnaeus, 1758) in southern Europe (Lepidoptera: Pieridae). *Nota Lepidopterologica* **11**: 7-20.
- Geiger H, Shapiro AM. 1992.** Genetics, systematics and evolution of Holarctic *Pieris napi* species group populations (Lepidoptera, Pieridae). *Zeitschrift für Zoologische Systematik und Evolutionsforschung* **30**: 100-122.
- Gere G. 1995.** Einfluß der Art *Colias erate* Esper (Lepidoptera, Pieridae) auf verwandte Spezies. *Opuscula Zoologica Budapest* **27-28**: 49-52.
- Gómez Bustillo MR, Fernández Rubio F. 1974.** *Las Mariposas de la Península Ibérica. Rhopaloceros*. Madrid: ICONA, Ministerio de Agricultura.
- Goodman OR, Turner HJ, Pierce FN. 1925.** A supposed hybrid *Polyommatus (coridon-hylas)*. *Entomological Record and Journal of Variation* **37**: 21-25.
- Groß F.J. 1957.** Ein vermutlicher Bastard zwischen *Coenonympha arcania* L. und *C. hero* L. *Entomologische Zeitschrift* **66**: 169-170.
- Guillaumin M, Descimon H. 1976.** La notion d'espèce chez les lépidoptères. In: Bocquet C, ed. *Les Problèmes de l'Espèce dans le Règne Animal*. Paris: Société zoologique de France, 129-201.
- Herrmann O. 1926.** Ein Hybrid von *Lyc. corydon* x *Lyc. icarus*? *Internationale Entomologische Zeitschrift* **19**: 380-381.
- Heslop Harrison JW. 1951.** A hybrid between *Pieris napi* male and *P. rapae* female (Lep., Pieridae). *Entomologist* **84**: 99-102.
- Higgins LG. 1969.** Observations sur les *Melanargia* dans le midi de la France. *Alexanor* **6**: 85-90.
- Higgins LG, Riley ND. 1970.** *A Field Guide to the Butterflies of Britain and Europe*. London: Collins.
- Kaisila J. 1950.** Über die vermuteten Bastarde zwischen *Colias hecla sulitelma* Aur. und *C. nastes werdandi* Zett. (Lep., Pieridae). *Annales Entomologici Fennici* **16**: 112-121.

- Klemann P. 1930.** Ein neuer Tagfalterhybrid. *Internationale Entomologische Zeitschrift* **23**: 438-442.
- Lux C. 1987.** Répartition en vertical des lépidoptères dans les Alpes Maritimes. Observation de phénomènes naturels. *Bulletin de la Société des Sciences Naturelles* **55**: 3-8.
- Mecke C. 1923.** *Colias myrmidone* und Hybrid *myrmidone* m. x *hyale* f. in Ostpreussen. *Internationale Entomologische Zeitschrift* **16**: 173-175.
- Peter W. 1928.** Zwitter von *Lycaena argus* L. und *Lycaena icarus* Rott. *Internationale Entomologische Zeitschrift* **22**: 139-140.
- Plantrou J. 1951.** Capture d'un hybride *hero-arcania*. *Revue Française de Lépidoptérologie* **13**: 129.
- Porter AH, Geiger H. 1995.** Limitations to the inference of gene flow at regional geographic scales - an example from the *Pieris napi* group (Lepidoptera: Pieridae) in Europe. *Biological Journal of the Linnean Society* **54**: 329-348.
- Porter AH, Wenger R, Geiger H, Scholl A, Shapiro AM. 1997.** The *Pontia daplidice-edusa* hybrid zone in northwestern Italy. *Evolution* **51**: 1561-1573.
- Rebel H. 1920.** *Lycaena* hybr. *meledamon* m. (*Lycaena meleager* Esp. m. x *Lyc. damon* Schiff. f.). *Verhandlungen der Zoologisch-Botanischen Gesellschaft. Wien* **70**: (75)-(77).
- Rebel H. 1929a.** *Lycaena* hybr. *bion* Rbl. m. (*L. damon* S.V. m. X *L. icarus* Rott f.). *Verhandlungen der Zoologisch-Botanischen Gesellschaft. Wien* **79**: (38)-(40).
- Rebel H. 1929b.** *Lycaena* hybr. *corydamon*-m. (*L. corydon* Pod. m. x *L. damon* Schiff. f.). *Verhandlungen der Zoologisch-Botanischen Gesellschaft. Wien* **79**: (33)-(36).
- Röber J. 1909.** Gattung: *Colias* F. In: Seitz A, ed. *Die Groß-Schmetterlinge der Erde*. Stuttgart: Verlag des Seitzschen Werkes (Alfred Kernen), 62-70.
- Rougeot PC. 1977.** Un hybride presumé d'*Anthocaris* grec. *Alexanor* **10**: 50.
- Ryszka H. 1949.** Ein *Colias* Hybride *ex ovo*. (*Colias hyale* L. m. x *Col. croceus* Fourc. f.). *Wiener Entomologische Rundschau* **1**: 5-7.
- Schurian K, Hoffmann P. 1975.** Ein neuer *Lycaeniden*-Hybrid von *Agrodiaetus damon* Schiff. *A. ripartii* Frt. (Lep., *Lycaenidae*). *Atalanta* **6**: 227-231.
- Tavoillot C. 1967.** Un hybride probable entre *Melanargia russiae* et *M. lachesis*. *Alexanor* **5**: 19-25.
- Tolman T, Lewington R. 1997.** *Butterflies of Britain and Europe*. London: HarperCollins.
- Wagner S. 1984.** *Melanargia lachesis* Hübner 1790, et-elle une espèce différente de *Melanargia galathea* Linnaeus 1758, oui ou non? *Nota Lepidopterologica* **7**: 375-386.

- Wenger R, Geiger H, Scholl A. 1993.** Genetic analysis of a hybrid zone between *P. daplidice* (Linnaeus, 1758) and *P. edusa* (Fabricius, 1777) (Lepidoptera: Pieridae) in Northern Italy. *Proceedings of the Fourth Congress of the European Society for Evolutionary Biology Abstracts*: 488.
- Wynhoff I. 1998.** The recent distribution of *Maculinea* species. *Journal of Insect Conservation* **2**: 15-27.